arXiv.org > stat > arXiv:1106.3921

Search or Article-id

(Help | Advanced search)

All papers



# Download:

- PDF
- Other formats

# Current browse context:

stat.ML

< prev | next > new | recent | 1106

#### Change to browse by:

q-fin q-fin.RM q-fin.ST stat.ME

#### References & Citations

NASA ADS

#### Bookmark(what is this?)











Statistics > Machine Learning

# **Dynamic Large Spatial Covariance Matrix Estimation in Application to** Semiparametric Model **Construction via Variable** Clustering: the SCE approach

## Song Song

(Submitted on 20 Jun 2011 (v1), last revised 23 Jun 2011 (this version, v2))

To better understand the spatial structure of large panels of economic and financial time series and provide a guideline for constructing semiparametric models, this paper first considers estimating a large spatial covariance matrix of the generalized \$m\$-dependent and \$\beta\$-mixing time series (with \$J\$ variables and \$T\$ observations) by hard thresholding regularization as long as  ${\{\log J \setminus cx^*(\ct)\}}/{T} = Co(1)$ \$ (the former scheme with some time dependence measure  $\c^*(\c)$  or  $\d J/\{T\} = \c^1$  (the latter scheme with some upper bounded mixing coefficient). We quantify the interplay between the estimators' consistency rate and the time dependence level, discuss an intuitive resampling scheme for threshold selection, and also prove a general cross-validation result justifying this. Given a consistently estimated covariance (correlation) matrix, by utilizing its natural links with graphical models and semiparametrics, after "screening" the (explanatory) variables, we implement a novel forward (and backward) label permutation procedure to cluster the "relevant" variables and construct the corresponding semiparametric model, which is further estimated by the groupwise dimension reduction method with sign constraints. We call this the SCE (screen - cluster estimate) approach for modeling high dimensional data with complex spatial structure. Finally we apply this method to study the spatial structure of large panels of economic and financial time series and find the proper semiparametric structure for estimating the consumer price index (CPI) to illustrate its superiority over the linear models.

Subjects: **Machine Learning (stat.ML)**; Risk Management (q-fin.RM);

Statistical Finance (q-fin.ST); Methodology (stat.ME)

Cite as: arXiv:1106.3921 [stat.ML]

(or arXiv:1106.3921v2 [stat.ML] for this version)

### Submission history

From: Song Song [view email]

[v1] Mon, 20 Jun 2011 14:35:52 GMT (1500kb,D) [v2] Thu, 23 Jun 2011 06:08:31 GMT (643kb,D)

Which authors of this paper are endorsers?

Link back to: arXiv, form interface, contact.