



Mathematical Physics

The path integral representation kernel of evolution operator in Merton-Garman model

L. F. Blazhyevskiy, V. S. Yanishevsky

(Submitted on 25 Jun 2011)

In the framework of path integral the evolution operator kernel for the Merton-Garman Hamiltonian is constructed. Based on this kernel option formula is obtained, which generalizes the well-known Black-Scholes result. Possible approximation numerical schemes for path integral calculations are proposed.

Comments: 16 pages

Subjects: **Mathematical Physics (math-ph)**; Pricing of Securities (q-fin.PR)

Journal reference: Condens. Matter Phys., 2011, vol. 14, No. 2, 23001:1-16

DOI: [10.5488/CMP.14.23001](https://doi.org/10.5488/CMP.14.23001)

Cite as: [arXiv:1106.5143](https://arxiv.org/abs/1106.5143) [math-ph]

(or [arXiv:1106.5143v1](https://arxiv.org/abs/1106.5143v1) [math-ph] for this version)

Submission history

From: CMPj [[view email](#)]

[v1] Sat, 25 Jun 2011 15:21:26 GMT (16kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math-ph

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[math](#)

[q-fin](#)

[q-fin.PR](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

