

Journal of Athletic Training

Home For Journal For Authors For Reviewers For Readers For Subscribers For Students Help

Home > [Journal of Athletic Training](#) > [May 2009](#) > Association of Quadriceps and Hamstrings Cocontraction Patterns With K...

[Advanced Search](#)

National Athletic Trainers' Association Links

- [NATA Home](#)
- [Online Manuscript Submission and Review](#)
- [Advertising](#)
- [Facts & Figures](#)
- [Editor-in-Chief](#)
- [Journal Editors](#)
- [Editorial Board](#)
- [NATA Position Statements](#)
- [PubMed Central](#)
- [Search PubMed](#)
- [Contact Us](#)

◀ [Previous Article](#) [Volume 44, Issue 3 \(May 2009\)](#) [Next Article](#) ▶

 [Add to Favorites](#)  [Share Article](#)  [Export Citations](#)

 [Track Citations](#)  [Permissions](#)

[Full-text](#)

[PDF](#)

Article Citation:

Riann M. Palmieri-Smith, Scott G. McLean, James A. Ashton-Miller, Edward M. Wojtys (2009) Association of Quadriceps and Hamstrings Cocontraction Patterns With Knee Joint Loading. *Journal of Athletic Training*: May 2009, Vol. 44, No. 3, pp. 256-263.

doi: 10.4085/1062-6050-44.3.256

Original Research

Association of Quadriceps and Hamstrings Cocontraction Patterns With Knee Joint Loading

Riann M. Palmieri-Smith, PhD, ATC, Scott G. McLean, PhD, James A. Ashton-Miller, PhD, and Edward M. Wojtys, MD

University of Michigan, Ann Arbor, MI

Abstract

Context: Sex differences in neuromuscular control of the lower extremity have been identified as a potential cause for the greater incidence of anterior cruciate ligament (ACL) injuries in female athletes compared with male athletes. Women tend to land in greater knee valgus with higher abduction loads than men. Because knee abduction loads increase ACL strain, the inability to minimize these loads may lead to ACL failure.

Objective: To investigate the activation patterns of the quadriceps and hamstrings muscles with respect to the peak knee abduction moment.

Design: Cross-sectional study.

Setting: Neuromuscular research laboratory.

Patients or Other Participants: Twenty-one recreationally active adults (11 women, 10 men).

Main Outcome Measure(s): Volunteers performed 3 trials of a 100-cm forward hop. During the hop task, we recorded surface electromyographic data from the medial and lateral hamstrings and quadriceps and recorded lower extremity kinematics and kinetics. Lateral and medial quadriceps-to-hamstrings (Q : H) cocontraction indices, the ratio of medial-to-lateral Q : H cocontraction, normalized root mean square electromyographic data for medial and lateral quadriceps and hamstrings, and peak knee abduction moment were calculated and used in data analyses.

Results: Overall cocontraction was lower in women than in men, whereas activation was lower in the medial than in the lateral musculature in both sexes ($P < .05$). The medial Q : H cocontraction index ($R^2 = 0.792$) accounted for a significant portion of the variance in the peak knee abduction moment in women ($P = .001$). Women demonstrated less activation in the vastus medialis than in the vastus lateralis ($P = .49$) and less activation in the medial hamstrings than in the

Volume 44, Issue 3
(May 2009)

< [Previous](#) [Next](#) >



[Current Issue](#)
[Available Issues](#)

Journal Information

Print ISSN 1062-6050

eISSN 1938-162X

Frequency Bimonthly:

January/February
March/April
May/June
July/August
September/October
November/December

Register for a Profile

Not Yet [Registered?](#)

Benefits of Registration Include:

- A Unique User Profile that will allow you to manage your current subscriptions (including online access)
- The ability to create favorites lists down to the article level
- The ability to customize email alerts to receive specific notifications about the topics you care most about and special offers

[Register Now!](#)

Related Articles

Articles Citing this Article

[Google Scholar](#)

Search for Other Articles By Author

- ☞ Riann M. Palmieri-Smith
- ☞ Scott G. McLean
- ☞ James A. Ashton-Miller
- ☞ Edward M. Wojtys

Search in:

Athletic Training

lateral hamstrings ($P = .01$).

Conclusions: Medial-to-lateral Q : H cocontraction appears to be unbalanced in women, which may limit their ability to resist abduction loads. Because higher abduction loads increase strain on the ACL, restoring medial-to-lateral Q : H cocontraction balance in women may help reduce ACL injury risk.

Keywords: [neuromuscular system](#), [biomechanics](#), [landings](#), [anterior cruciate ligament](#), [coactivation](#)

Riann M. Palmieri-Smith, PhD, ATC, contributed to conception and design; acquisition and analysis and interpretation of the data; and drafting, critical revision, and final approval of the article. Scott G. McLean, PhD, contributed to analysis and interpretation of the data and drafting, critical revision, and final approval of the article. James A. Ashton-Miller, PhD, and Edward M. Wojtys, MD, contributed to conception and design and drafting, critical revision, and final approval of the article.

Riann M. Palmieri-Smith, PhD, ATC, Athletic Training, Movement Science, & Orthopaedics, 4745G CCRB, 401 Washtenaw Avenue, School of Kinesiology, University of Michigan, Ann Arbor, MI 48109, e-mail: riannp@umich.edu

[top](#) ▲