

Jordanian deformation of the open XXX-spin chain

P. P. Kulish, N. Manojlovic, Z. Nagy

(Submitted on 30 Nov 2009 (v1), last revised 18 Dec 2009 (this version, v2))

The general solution to the reflection equation associated with the Jordanian deformation of the $SL(2)$ invariant Yang R-matrix is found. The same K-matrix is obtained by the special scaling limit of the XXZ-model with general boundary conditions. The Hamiltonian with the boundary terms is explicitly derived according to the Sklyanin formalism. We discuss the structure of the spectrum of the deformed XXX-model and its dependence on the boundary conditions.

Comments: 13 pages; typos corrected

Subjects: **Exactly Solvable and Integrable Systems (nlin.SI)**; High Energy Physics - Theory (hep-th); Quantum Algebra (math.QA)

Cite as: [arXiv:0911.5592v2](#) [nlin.SI]

Submission history

From: Nenad Manojlovic [[view email](#)]

[v1] Mon, 30 Nov 2009 09:38:56 GMT (13kb)

[v2] Fri, 18 Dec 2009 15:35:04 GMT (13kb)

Which authors of this paper are endorsers?

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nlin.SI

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [0911](#)

Change to browse by:

[hep-th](#)

[math](#)

[math.QA](#)

[nlin](#)

References & Citations

- [CiteBase](#)

Bookmark([what is this?](#))

[CiteULike logo](#)

[Connotea logo](#)

[BibSonomy logo](#)

[Mendeley logo](#)

[Facebook logo](#)

[del.icio.us logo](#)

[Digg logo](#)

[Reddit logo](#)