

Comparison of Dualizing Complexes

Changlong Zhong

(Submitted on 12 Nov 2010)

We prove that there is a map from Bloch's cycle complex to Kato's complex of Milnor K-theory, which induces a quasi-isomorphism from \mathcal{E} 's sheafified cycle complex to the Gersten complex of logarithmic de Rham--Witt sheaves. Next we show that the truncation of Bloch's cycle complex at -3 is quasi-isomorphic to Spiess' dualizing complex.

Comments: 23 pages

Subjects: **Algebraic Geometry (math.AG)**; K-Theory and Homology (math.KT); Number Theory (math.NT)

MSC classes: 14F20, 14C25, 11G25, 19E15

Cite as: [arXiv:1011.2826v1](https://arxiv.org/abs/1011.2826v1) [math.AG]

Submission history

From: Changlong Zhong [[view email](#)]

[v1] Fri, 12 Nov 2010 04:52:18 GMT (17kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AG

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1011](#)

Change to browse by:

[math](#)

[math.KT](#)

[math.NT](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

