

# Some two-dimensional finite energy percolation processes

Olle Hägström, Péter Mester

(Submitted on 12 Nov 2010)

Some examples of translation invariant site percolation processes on the  $\mathbb{Z}^2$  lattice are constructed, the most far-reaching example being one that satisfies uniform finite energy (meaning that the probability that a site is open given the status of all others is bounded away from 0 and 1) and exhibits a.s. the coexistence of an infinite open cluster and an infinite closed cluster. Essentially the same example shows that coexistence is possible between an infinite open cluster and an infinite closed cluster that are both robust under i.i.d. thinning.

Comments: 14 pages

Subjects: **Probability (math.PR)**

Journal reference: Electr. Comm. Probab. 14, 42-54. 2009

Cite as: [arXiv:1011.2872v1](#) [math.PR]

## Submission history

From: Péter Mester [[view email](#)]

[v1] Fri, 12 Nov 2010 10:20:27 GMT (16kb)

*Which authors of this paper are endorsers?*

Link back to: [arXiv](#), [form interface](#), [contact](#).

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

**math.PR**

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1011](#)

Change to browse by:

[math](#)

## References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

