

Translation invariant equations and the method of Sanders

Thomas Bloom

(Submitted on 6 Jul 2011 (v1), last revised 30 Jan 2012 (this version, v2))

We extend the recent improvement of Roth's theorem on three term arithmetic progressions by Sanders to obtain similar results for the problem of locating non-trivial solutions to translation invariant linear equations in many variables in both $\mathbb{Z}/N\mathbb{Z}$ and $\mathbb{F}_q[t]$.

Subjects: **Number Theory (math.NT)**; Combinatorics (math.CO)

DOI: [10.1112/blms/bds045](https://doi.org/10.1112/blms/bds045)

Cite as: [arXiv:1107.1110](https://arxiv.org/abs/1107.1110) [math.NT]

(or [arXiv:1107.1110v2](https://arxiv.org/abs/1107.1110v2) [math.NT] for this version)

Submission history

From: Thomas Bloom [[view email](#)]

[v1] Wed, 6 Jul 2011 12:43:20 GMT (17kb)

[v2] Mon, 30 Jan 2012 11:14:55 GMT (18kb,D)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [Other formats](#)

Current browse context:

math.NT

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math.CO](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

