

# Young's seminormal form and simple modules for $S_n$ in characteristic $p$

Steen Ryom-Hansen

(Submitted on 15 Jul 2011 (v1), last revised 5 Sep 2012 (this version, v3))

We realize the integral Specht modules for the symmetric group  $S_n$  as induced modules from the subalgebra of the group algebra generated by the Jucys-Murphy elements. We deduce from this that the simple modules for  $FS_n$  are generated by reductions modulo  $p$  of the corresponding Jucys-Murphy idempotents.

Comments: Published version Algebras and Representation Theory 2012

Subjects: **Representation Theory (math.RT)**; Quantum Algebra (math.QA)

Cite as: **arXiv:1107.3076 [math.RT]**

(or **arXiv:1107.3076v3 [math.RT]** for this version)

## Submission history

From: Steen Ryom-Hansen [[view email](#)]

[v1] Fri, 15 Jul 2011 14:33:11 GMT (30kb)

[v2] Fri, 20 Jan 2012 15:53:30 GMT (30kb)

[v3] Wed, 5 Sep 2012 15:07:50 GMT (33kb)

*[Which authors of this paper are endorsers?](#)*

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.RT

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math.QA](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))



Science  
WISE