



Aczélian n-ary semigroups

[Miguel Couceiro](#), [Jean-Luc Marichal](#)

(Submitted on 25 Jul 2011 ([v1](#)), last revised 3 Aug 2012 (this version, [v2](#)))

We show that the real continuous, symmetric, and cancellative n-ary semigroups are topologically order-isomorphic to additive real n-ary semigroups. The binary case (n=2) was originally proved by Acz'el (1949); there symmetry was redundant.

Subjects: **Rings and Algebras (math.RA)**; Functional Analysis (math.FA)

MSC classes: 20N15, 39B22 (Primary) 20M14 (Secondary)

Journal reference: Semigroup Forum 85 (1) (2012) 81-90

DOI: [10.1007/s00233-011-9354-2](https://doi.org/10.1007/s00233-011-9354-2)

Cite as: [arXiv:1107.4989](#) [math.RA]
(or [arXiv:1107.4989v2](#) [math.RA] for this version)

Submission history

From: Jean-Luc Marichal [[view email](#)]

[\[v1\]](#) Mon, 25 Jul 2011 16:09:27 GMT (10kb)

[\[v2\]](#) Fri, 3 Aug 2012 11:12:10 GMT (10kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.RA

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math.FA](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

