

Nonlinear Two-Dimensional Green's Function in Smectics

E A. Brener, V. I. Marchenko, D. Pilipenko

(Submitted on 21 Mar 2010)

The problem of the strain of smectics subjected to a force distributed over a line in the basal plane has been solved.

Subjects: **Materials Science (cond-mat.mtrl-sci)**; Soft Condensed Matter (cond-mat.soft); Pattern Formation and Solitons (nlin.PS)

Journal reference: Pis'ma Zh. Eksp. Teor. Fiz. vol. 90, 761 (2009)

Cite as: [arXiv:1003.3995v1](#) [cond-mat.mtrl-sci]

Submission history

From: Denis Pilipenko [[view email](#)]

[v1] Sun, 21 Mar 2010 13:41:03 GMT (8kb)

Which authors of this paper are endorsers?

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

cond-mat.mtrl-sci

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1003](#)

Change to browse by:

[cond-mat](#)

[cond-mat.soft](#)

[nlin](#)

[nlin.PS](#)

References & Citations

- [CiteBase](#)

Bookmark (what is this?)

 [CiteULike logo](#)

 [Connotea logo](#)

 [BibSonomy logo](#)

 [Mendeley logo](#)

 [Facebook logo](#)

 [del.icio.us logo](#)

 [Digg logo](#)

 [Reddit logo](#)