

Darboux transformation for two component derivative nonlinear Schrödinger equation

Liming Ling, Q. P. Liu

(Submitted on 25 Nov 2009 (v1), last revised 9 Dec 2009 (this version, v2))

In this paper, we consider the two component derivative nonlinear Schrödinger equation and present a simple Darboux transformation for it. By iterating this Darboux transformation, we construct a compact representation for the N -soliton solutions.

Comments: 12 pages, 2 figures

Subjects: **Exactly Solvable and Integrable Systems (nlin.SI)**

Cite as: **arXiv:0911.4802v2 [nlin.SI]**

Submission history

From: Q. P. Liu [[view email](#)]

[v1] Wed, 25 Nov 2009 10:26:42 GMT (206kb)

[v2] Wed, 9 Dec 2009 07:50:07 GMT (449kb)

Which authors of this paper are endorsers?

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nlin.SI

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [0911](#)

Change to browse by:

[nlin](#)

References & Citations

- [CiteBase](#)

Bookmark([what is this?](#))

[CiteULike logo](#)

[Connotea logo](#)

[BibSonomy logo](#)

[Mendeley logo](#)

[Facebook logo](#)

[del.icio.us logo](#)

[Digg logo](#)

[Reddit logo](#)