

# Index character associated to the projective Dirac operator

Makoto Yamashita

(Submitted on 15 Jul 2011)

We calculate the equivariant index formula for an infinite dimensional Clifford module canonically associated to any Riemannian manifold. It encompasses the fractional index formula of the projective Dirac operator by Mathai--Melrose--Singer.

Comments: 8 pages

Subjects: **K-Theory and Homology (math.KT)**; Differential Geometry (math.DG)

MSC classes: 58J22 (primary) 35K05 (secondary)

Cite as: [arXiv:1107.3007](https://arxiv.org/abs/1107.3007) [math.KT]

(or [arXiv:1107.3007v1](https://arxiv.org/abs/1107.3007v1) [math.KT] for this version)

## Submission history

From: Makoto Yamashita [[view email](#)]

[v1] Fri, 15 Jul 2011 08:29:17 GMT (11kb)

*Which authors of this paper are endorsers?*

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

## Current browse context:

math.KT

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

## Change to browse by:

[math](#)

[math.DG](#)

## References & Citations

- [NASA ADS](#)

## Bookmark (what is this?)



Science  
WISE