



Mathematics > Algebraic Geometry

Harmonic analysis and the Riemann-Roch theorem

D. V. Osipov, A. N. Parshin

(Submitted on 2 Jul 2011 (v1), last revised 21 Oct 2011 (this version, v2))

This paper is a continuation of papers: [arXiv:0707.1766](#) [math.AG] and [arXiv:0912.1577](#) [math.AG]. Using the two-dimensional Poisson formulas from these papers and two-dimensional adelic theory we obtain the Riemann-Roch formula on a projective smooth algebraic surface over a finite field.

Comments: 7 pages; to appear in Doklady Mathematics

Subjects: **Algebraic Geometry (math.AG)**; Number Theory (math.NT)

Cite as: [arXiv:1107.0408v2](#) [math.AG]

Submission history

From: Denis V. Osipov [[view email](#)]

[v1] Sat, 2 Jul 2011 19:58:36 GMT (8kb)

[v2] Fri, 21 Oct 2011 18:01:26 GMT (9kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AG

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math.NT](#)

References & Citations

- [NASA ADS](#)

Bookmark (what is this?)

