



Toric Hirzebruch-Riemann-Roch via Ishida's theorem on the Todd genus

Hal Schenck

(Submitted on 3 Jul 2011 (v1), last revised 24 Aug 2011 (this version, v2))

We give a simple proof of the Hirzebruch-Riemann-Roch theorem for smooth complete toric varieties, based on Ishida's result that the Todd genus of a smooth complete toric variety is one.

Comments: 3 pages, v2 title change, references updated

Subjects: **Algebraic Geometry (math.AG)**; Commutative Algebra (math.AC)

MSC classes: 14M25, 14C40

Cite as: **arXiv:1107.0489 [math.AG]**

(or **arXiv:1107.0489v2 [math.AG]** for this version)

Submission history

From: Henry K. Schenck [[view email](#)]

[v1] Sun, 3 Jul 2011 19:49:19 GMT (4kb)

[v2] Wed, 24 Aug 2011 13:56:58 GMT (4kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AG

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math.AC](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

