

# Sources of log canonical centers

János Kollár (Princeton Univ)

(Submitted on 14 Jul 2011 (v1), last revised 14 Nov 2012 (this version, v3))

We define the "source" and the "spring" of a log canonical center and use them to solve several problems in higher-codimension adjunction. The main application is to the construction of semi log canonical pairs.

Version 2: References updated. Version 3: Many small changes.

Subjects: **Algebraic Geometry (math.AG)**

MSC classes: 14J17, 14E30, 14J15

Cite as: **arXiv:1107.2863 [math.AG]**

(or **arXiv:1107.2863v3 [math.AG]** for this version)

## Submission history

From: János Kollár [[view email](#)]

[v1] Thu, 14 Jul 2011 16:20:41 GMT (24kb)

[v2] Thu, 21 Jul 2011 19:28:12 GMT (24kb)

[v3] Wed, 14 Nov 2012 17:42:25 GMT (20kb)

[Which authors of this paper are endorsers?](#)

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

## Current browse context:

math.AG

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

## Change to browse by:

[math](#)

## References & Citations

- [NASA ADS](#)

## 1 blog link (what is this?)

## Bookmark (what is this?)



Science  
WISE