

On the unramified Brauer group of a homogeneous space

Mikhail Borovoi

(Submitted on 5 Jun 2012 (v1), last revised 10 Sep 2012 (this version, v2))

We give a new proof of the theorem stating that for any connected linear algebraic group G over an algebraically closed field k of characteristic 0 and for any closed connected subgroup H of G , the unramified Brauer group of G/H vanishes.

Comments: 4 pages; shortened, misprints corrected

Subjects: **Algebraic Geometry (math.AG)**

MSC classes: 14F22 (Primary) 14M17, 14L10 (Secondary)

Cite as: **arXiv:1206.1023 [math.AG]**

(or **arXiv:1206.1023v2 [math.AG]** for this version)

Submission history

From: Mikhail Borovoi [[view email](#)]

[v1] Tue, 5 Jun 2012 19:05:54 GMT (6kb)

[v2] Mon, 10 Sep 2012 16:22:25 GMT (5kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AG

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1206](#)

Change to browse by:

[math](#)

References & Citations

- [NASA ADS](#)

Bookmark (what is this?)



Science
WISE