Cornell University Library

We gratefully acknowledge support from the Simons Foundation and member institutions

arXiv.org > math > arXiv:1107.0106

Search or Article-id

(Help | Advanced search)

All papers | Go!

Mathematics > Differential Geometry

Calabi-Yau Problem for Legendrian curves in C^3 and applications

Francisco Martin, Masaaki Umehara, Kotaro Yamada

(Submitted on 1 Jul 2011 (v1), last revised 23 May 2012 (this version, v2))

We construct a complete, bounded Legendrian immersion in C^3. As direct applications of it, we show the first examples of a weakly complete bounded flat front in hyperbolic 3-space, a weakly complete bounded flat front in de Sitter 3-space, and a weakly complete bounded improper affine front in R^3.

Comments:	After the submission, the authors found that the proof of the main theorem contains a serious error and the main theorem turns to be an open problem. ArXiv:1205.4779 is a new paper showing the existence of a weakly complete flat surface (with singularities) in hyperbolic 3-space such that the images of the hyperbolic Gauss maps are bounded, which plays a role of a recovery of the present paper
Subjects:	**Differential Geometry (math.DG)**
MSC classes:	53D10, 53A10, 53A1, 53A35
Cite as:	**arXiv:1107.0106** [math.DG]
	(or **arXiv:1107.0106v2** [math.DG] for this version)

Submission history

From: Kotaro Yamada [view email]

[v1] Fri, 1 Jul 2011 05:28:42 GMT (17kb)

[v2] Wed, 23 May 2012 10:14:59 GMT (17kb)

Which authors of this paper are endorsers?