arXiv.org > math > arXiv:1107.5621

Search or Article-id

(Help | Advanced search)

All papers



Mathematics > Geometric Topology

A tour of bordered Floer theory

Robert Lipshitz, Peter S. Ozsvath, Dylan P. Thurston

(Submitted on 28 Jul 2011)

Heegaard Floer theory is a kind of topological quantum field theory, assigning graded groups to closed, connected, oriented 3-manifolds and group homomorphisms to smooth, oriented 4-dimensional cobordisms. Bordered Heegaard Floer homology is an extension of Heegaard Floer homology to 3-manifolds with boundary, with extended-TQFT-type gluing properties. In this survey, we explain the formal structure and construction of bordered Floer homology and sketch how it can be used to compute some aspects of Heegaard Floer theory.

Comments: 13 pages, 7 figures

Subjects: Geometric Topology (math.GT); Quantum Algebra

(math.QA); Symplectic Geometry (math.SG)

MSC classes: 57R (Primary) 53D40 (Secondary) Journal reference: Proc.Nat.Acad.Sci.108:8085-8092,2011

DOI: 10.1073/pnas.1019060108 Cite as: arXiv:1107.5621v1 [math.GT]

Submission history

From: Dylan Thurston [view email]

[v1] Thu, 28 Jul 2011 04:45:01 GMT (121kb,D)

Which authors of this paper are endorsers?

Link back to: arXiv, form interface, contact.

Download:

- PDF
- Other formats

Current browse context:

math.GT

< prev | next > new | recent | 1107

Change to browse by:

math

math.QA math.SG

References & Citations

NASA ADS

2 blog links(what is this?)

Bookmark(what is this?)











