

Bäcklund Transformations for the Trigonometric Gaudin Magnet

Orlando Ragnisco, Federico Zullo

(Submitted on 12 Dec 2009 (v1), last revised 29 Jan 2010 (this version, v2))

We construct a Backlund transformation for the trigonometric classical Gaudin magnet starting from the Lax representation of the model. The Darboux dressing matrix obtained depends just on one set of variables because of the so-called spectrality property introduced by E. Sklyanin and V. Kuznetsov. In the end we mention some possibly interesting open problems.

Comments: contribution to the Proc. of "Integrable Systems and Quantum Symmetries 2009", Prague, June 18-20, 2009

Subjects: **Exactly Solvable and Integrable Systems (nlin.SI)**

Journal reference: SIGMA 6 (2010), 012, 6 pages

DOI: [10.3842/SIGMA.2010.012](https://doi.org/10.3842/SIGMA.2010.012)

Cite as: [arXiv:0912.2456v2](https://arxiv.org/abs/0912.2456v2) [nlin.SI]

Submission history

From: Orlando Ragnisco [[view email](#)]

[v1] Sat, 12 Dec 2009 22:12:41 GMT (8kb)

[v2] Fri, 29 Jan 2010 06:32:21 GMT (10kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nlin.SI

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [0912](#)

Change to browse by:

[nlin](#)

References & Citations

- [CiteBase](#)

Bookmark([what is this?](#))

[CiteULike logo](#)

[Connotea logo](#)

[BibSonomy logo](#)

[Mendeley logo](#)

[Facebook logo](#)

[del.icio.us logo](#)

[Digg logo](#)

[Reddit logo](#)