

Vertex operator for the non-autonomous ultradiscrete KP equation

[Yoichi Nakata](#)

(Submitted on 22 Dec 2009)

We propose an ultradiscrete analogue of the vertex operator in the case of the ultradiscrete KP equation--several other ultradiscrete equations--which maps N-soliton solutions to N+1-soliton ones.

Comments: 9 pages

Subjects: **Exactly Solvable and Integrable Systems (nlin.SI)**

Cite as: [arXiv:0912.4302v1](#) [nlin.SI]

Submission history

From: Yoichi Nakata [[view email](#)]

[v1] Tue, 22 Dec 2009 00:47:05 GMT (20kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nlin.SI

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [0912](#)

Change to browse by:

[nlin](#)

References & Citations

- [CiteBase](#)

Bookmark([what is this?](#))

[CiteULike](#) logo

[Connotea](#) logo

[BibSonomy](#) logo

[Mendeley](#) logo

[Facebook](#) logo

[del.icio.us](#) logo

[Digg](#) logo

[Reddit](#) logo