



# Strong approximation of almost periodic functions

[Włodzimierz Łenski](#), [Bogdan Szal](#)

(Submitted on 13 Apr 2012)

We consider summability methods generated by the class GM(2b). We generalize some related results of P. Pych-Taberska [Studia Math. XCVI (1990), 91-103] on strong approximation of almost periodic functions by their Fourier series and S. M. Mazhar and V. Totik [J. Approx. Theory, 60(1990), 174-182] on approximation of periodic functions by matrix means of their Fourier series.

Subjects: **Classical Analysis and ODEs (math.CA)**

Cite as: **arXiv:1204.2938 [math.CA]**

(or **arXiv:1204.2938v1 [math.CA]** for this version)

## Submission history

From: Bogdan Szal [[view email](#)]

[v1] Fri, 13 Apr 2012 10:06:05 GMT (8kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.CA

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1204](#)

Change to browse by:

[math](#)

## References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

