



Mathematics > Analysis of PDEs

Resonant uniqueness of radial semiclassical Schrodinger operators

Kiril Datchev, Hamid Hezari

(Submitted on 5 Jul 2011)

We prove that radial, monotonic, superexponentially decaying potentials in \mathbb{R}^n , n greater than or equal to 1 odd, are determined by the resonances of the associated semiclassical Schrodinger operator among all superexponentially decaying potentials in \mathbb{R}^n .

Subjects: **Analysis of PDEs (math.AP)**; Mathematical Physics (math-ph); Spectral Theory (math.SP)

Cite as: [arXiv:1107.0960v1](https://arxiv.org/abs/1107.0960v1) [math.AP]

Submission history

From: Kiril Datchev [[view email](#)]

[v1] Tue, 5 Jul 2011 19:46:34 GMT (8kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AP

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math-ph](#)

[math.SP](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

