



Extending cutoff resolvent estimates via propagation of singularities

[Kiril Datchev](#)

(Submitted on 14 Jul 2011 ([v1](#)), last revised 22 Jun 2012 (this version, [v2](#)))

We use a gluing method developed in joint work with Andr'as Vasy to show that polynomially bounded cutoff resolvent estimates at the real axis imply, up to a constant factor, the same estimates in a neighborhood of the real axis.

Comments: 5 pages

Subjects: **Analysis of PDEs (math.AP)**

Journal reference: Communications in Partial Differential Equations, Vol. 37, No. 8, pp. 1456--1461, 2012

Cite as: [arXiv:1107.2941](#) [math.AP]
(or [arXiv:1107.2941v2](#) [math.AP] for this version)

Submission history

From: Kiril Datchev [[view email](#)]

[\[v1\]](#) Thu, 14 Jul 2011 20:15:56 GMT (6kb)

[\[v2\]](#) Fri, 22 Jun 2012 21:32:04 GMT (6kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AP

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

