



A decay estimate for a wave equation with trapping and a complex potential

Lars Andersson, Pieter Blue, Jean-Philippe Nicolas

(Submitted on 22 Jul 2011)

In this brief note, we consider a wave equation that has both trapping and a complex potential. For this problem, we prove a uniform bound on the energy and a Morawetz (or integrated local energy decay) estimate. The equation is a model problem for certain scalar equations appearing in the Maxwell and linearised Einstein systems on the exterior of a rotating black hole.

Comments: 11 pages

Subjects: **Analysis of PDEs (math.AP)**; General Relativity and Quantum Cosmology (gr-qc)

MSC classes: 35Q75

Cite as: **arXiv:1107.4597 [math.AP]**

(or **arXiv:1107.4597v1 [math.AP]** for this version)

Submission history

From: Pieter Blue [[view email](#)]

[v1] Fri, 22 Jul 2011 19:26:06 GMT (12kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AP

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[gr-qc](#)

[math](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

