



A fast alternating projection method for complex frequency estimation

Fredrik Andersson, Marcus Carlsson, Per-Anders Ivert

(Submitted on 8 Jul 2011)

The problem of approximating a sampled function using sums of a fixed number of complex exponentials is considered. We use alternating projections between fixed rank matrices and Hankel matrices to obtain such an approximation. Convergence, convergence rates and error estimates for this technique are proven, and fast algorithms are developed. We compare the numerical results obtain with the MUSIC and ESPRIT methods.

Subjects: **Numerical Analysis (math.NA)**

Cite as: [arXiv:1107.2028](#) [math.NA]

(or [arXiv:1107.2028v1](#) [math.NA] for this version)

Submission history

From: Fredrik Andersson [[view email](#)]

[v1] Fri, 8 Jul 2011 14:16:14 GMT (54kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.NA

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

