



# Series Prediction based on Algebraic Approximants

Herbert H. H. Homeier

(Submitted on 12 Jul 2011)

It is described how the Hermite-Padé polynomials corresponding to an algebraic approximant for a power series may be used to predict coefficients of the power series that have not been used to compute the Hermite-Padé polynomials. A recursive algorithm is derived and some numerical examples are given.

Comments: 9 pages, ISRN Applied Mathematics, in press  
 Subjects: **Numerical Analysis (math.NA)**; Computational Physics (physics.comp-ph)  
 MSC classes: 41A25, 65D99  
 Journal reference: ISRN Applied Mathematics, vol. 2011, Article ID 958968, 7 pages, 2011  
 DOI: [10.5402/APPMATH 10.5402/2011/958968](https://doi.org/10.5402/APPMATH.10.5402/2011/958968)  
 Cite as: [arXiv:1107.2406](https://arxiv.org/abs/1107.2406) [math.NA]  
 (or [arXiv:1107.2406v1](https://arxiv.org/abs/1107.2406v1) [math.NA] for this version)

## Submission history

From: Herbert H. H. Homeier [[view email](#)]  
 [v1] Tue, 12 Jul 2011 20:02:05 GMT (6kb)

*[Which authors of this paper are endorsers?](#)*

Link back to: [arXiv](#), [form interface](#), [contact](#).

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.NA

< [prev](#) | [next](#) >

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[physics](#)

[physics.comp-ph](#)

## References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

