

# Numerical Methods for Multilattices

Assyr Abdulle, Ping Lin, Alexander V. Shapeev

(Submitted on 18 Jul 2011 (v1), last revised 10 Sep 2012 (this version, v2))

Among the efficient numerical methods based on atomistic models, the quasicontinuum (QC) method has attracted growing interest in recent years. The QC method was first developed for crystalline materials with Bravais lattice and was later extended to multilattices (Tadmor et al, 1999). Another existing numerical approach to modeling multilattices is homogenization. In the present paper we review the existing numerical methods for multilattices and propose another concurrent macro-to-micro method in the numerical homogenization framework. We give a unified mathematical formulation of the new and the existing methods and show their equivalence. We then consider extensions of the proposed method to time-dependent problems and to random materials.

Comments: 31 pages

Subjects: **Numerical Analysis (math.NA)**

MSC classes: 65N30, 70C20, 74G15, 74G65

Journal reference: Multiscale Model. Simul., Vol 10(3), pp. 696-726, 2012

DOI: [10.1137/110841163](https://doi.org/10.1137/110841163)

Cite as: [arXiv:1107.3462](https://arxiv.org/abs/1107.3462) [math.NA]

(or [arXiv:1107.3462v2](https://arxiv.org/abs/1107.3462v2) [math.NA] for this version)

## Submission history

From: Alexander V. Shapeev [[view email](#)]

[v1] Mon, 18 Jul 2011 15:07:57 GMT (919kb,D)

[v2] Mon, 10 Sep 2012 05:01:53 GMT (1028kb,D)

[Which authors of this paper are endorsers?](#)

## Download:

- [PDF](#)
- [Other formats](#)

## Current browse context:

math.NA

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1107](#)

## Change to browse by:

[math](#)

## References & Citations

- [NASA ADS](#)

## Bookmark (what is this?)

