



Mathematics > Statistics Theory

On Low-Dimensional Projections of High-Dimensional Distributions

Lutz Duembgen, Perla Zerial

(Submitted on 2 Jul 2011 (v1), last revised 12 Dec 2011 (this version, v3))

Let P be a probability distribution on q -dimensional space. The so-called Diaconis-Freedman effect means that for a fixed dimension $d \ll q$, most d -dimensional projections of P look like a scale mixture of spherically symmetric Gaussian distributions. The present paper provides necessary and sufficient conditions for this phenomenon in a suitable asymptotic framework with increasing dimension q . It turns out, that the conditions formulated by Diaconis and Freedman (1984) are not only sufficient but necessary as well. Moreover, letting \hat{P} be the empirical distribution of n independent random vectors with distribution P , we investigate the behavior of the empirical process $\sqrt{n}(\hat{P} - P)$ under random projections, conditional on \hat{P} .

Subjects: **Statistics Theory (math.ST)**
 MSC classes: 62E20, 62G20, 62H99
 Cite as: **arXiv:1107.0417v3 [math.ST]**

Submission history

From: Lutz Duembgen [view email]
[v1] Sat, 2 Jul 2011 22:11:14 GMT (12kb)
[v2] Thu, 27 Oct 2011 11:49:49 GMT (13kb)
[v3] Mon, 12 Dec 2011 09:43:27 GMT (13kb)

Which authors of this paper are endorsers?

Download:

- PDF
- PostScript
- Other formats

Current browse context:

math.ST

< prev | next >

new | recent | 1107

Change to browse by:

math
stat

References & Citations

- NASA ADS

Bookmark (what is this?)

