

Free boundary problems of the Euler equation: local well-posedness and hydrodynamical instabilities

Jalal Shatah
Courant Institute
New York University

Chongchun Zeng
School of Mathematics
Georgia Institute of Technology

Droplet in Vacuum (1F)

- Euler's equation (E): $v_t + v \cdot \nabla v = -\nabla p, \quad x \in \Omega_t$
 - $\Omega_t \subset \subset \mathbf{R}^n$: moving fluid domain
 - p : pressure
 - Incompressibility: $\nabla \cdot v = 0$
- Free boundary problem in vacuum:
 1. $v|_{\partial\Omega_t}$: velocity of the moving boundary
 2. $p|_{\partial\Omega_t} = \alpha\kappa$: surface tension, $\alpha \in [0, 1]$
 κ : mean curvature of $\partial\Omega_t \subset \mathbf{R}^n$

Interface Problem (2F)

- Ω_t^\pm : moving domains of two fluids
 - $S_t = \partial\Omega_t^+ = \partial\Omega_t^-$: moving interface
 - Euler's equation (E): $\rho_\pm(\partial_t v_\pm + v_\pm \cdot \nabla v_\pm) = -\nabla p_\pm, \quad x \in \Omega_t^\pm$
 - ρ_\pm : constant densities of two fluids
 - Incompressibility $\nabla \cdot v = 0$
- Free boundary problem:
 1. $v_+^\perp|_{S_t} = v_-^\perp|_{S_t}$: interface velocity
 2. $(p_+ - p_-)|_{S_t} = \kappa_+$: surface tension

- **Instabilities** in the absence of surface tension: ill-posedness
 - (1F): possible Rayleigh-Taylor instability. **[RT]**
 - (2F): Kelvin-Helmholtz instability. **[KH]**
- * (1F): if $\nabla \times v = 0$, then no **[RT]**
- * Surface tension: regularizing force, suppress **[RT]** or **[KH]**
- **Difficulties:**
 - * Fluid domain unknown
 - * Choice of coordinates: Lagrangian coordinates?

Reference I. Without surface tension

- Nalimov 74, Yoshihara 82: **(1F)**, irrotation, 2-d, small data → local well posedness
- Ebin 87: **(1F)**, ill-posedness with Rayleigh-Taylor instability
- Wu 97, 99 **(1F)** irrotational → local well-posedness; 06 **(2F)** irrotational
- Beyer & Gunther 00: **(1F)**, irrotational → linearization
- Christodoulou & Lindblad 00, Lindblad 03, ... : **(1F)**, energy estimates, local well-posedness
- Coutand & Shkoller 05, Zhang & Zhang 06: **(1F)**, local well-posedness

Reference II. With surface tension

- Yoshihara 83, Iguchi 01 (**1F**) 2d irrotational well-posedness
 - Ambrose 03: (**2F**) 2-d irrotational \rightarrow well-posedness
 - Beyer & Gunther 98, 05: (**1F**) irrotational \rightarrow well-posedness;
 - Masmoudi & Ambrose: 05: (**1F**) 2-d irrotational, 06: (**2F**) 3-d irrotational, well-posedness
 - Coutand & Shkoller & Cheng 05, 08: (**1F**), (**2F**) well-posedness
- Shinbrot 76, Nishida 79, Sulem & et. al. 81, Kano & Nishida 79, 86, Craig 85, Ogawa & Tani: 02, Schneider & Wayne 02, Lannes 04: 2-d, Schweizer: 05, Beale, Hou, & Lowengrub 93, Hou, Lowengrub, & Shelley 97, Christianson, Hur, & Staffinani
....

Irrotational and simply connected case: $\nabla \times v = 0$

- (a) $\exists \phi(t, x), x \in \Omega_t$, s.t. $\nabla \phi = v$
 - (b) $\nabla \cdot v = 0 \Rightarrow \Delta \phi = 0$
 - (c) $\frac{\partial \phi}{\partial N} = v \cdot N = v^\perp$ on $\partial \Omega_t$
-
- The problem is reduced to $\partial \Omega_t$
 - * Explicit pressure is avoided
 - * Nonlocal operators are involved

A first look at the pressure

$$-\Delta p = \partial_i v^j \partial_j v^i \quad \text{in } \Omega_t$$

- Fixed boundary + non-penetrating B.C. on $\partial\Omega$: $v \cdot N = 0$

$(E) \cdot N|_{\partial\Omega} \Rightarrow \nabla_N p|_{\partial\Omega} = -\nabla_v v \cdot N = v \cdot \nabla_v N \Rightarrow$ regularity: $\nabla p \sim v$

- **(1F)** w/o surface tension: $p = 0$ on $\partial\Omega_t \Rightarrow$

reg. of $\nabla p \leq$ (reg. of $\partial\Omega_t - \frac{1}{2}$) and (reg. of v)

- **(1F)** + surface tension: $p = \kappa$ on $\partial\Omega_t \Rightarrow$

reg. of $\nabla p \leq$ (reg. of $\partial\Omega_t - \frac{5}{2}$) and (reg. of v)

- Surface tension vs. No surface tension
 - (a) Physically, a force regularizing $\partial\Omega_t$
 - (b) Lower regularity of ∇p
- Regularity of $\partial\Omega_t$: best described by Lagrangian coordinates?
 - * Lagrangian coordinates \rightarrow reg. $\partial\Omega_t = \text{reg. } v - \frac{1}{2}$
 - * How to close the regularity count?

conservation of energy

$$v_t + v \cdot \nabla v = -\nabla p, \quad x \in \Omega_t \quad (\text{E})$$

* $v|_{\partial\Omega_t}$: boundary velocity, $p|_{\partial\Omega_t} = \kappa$: mean curvature of $\partial\Omega_t$

- $\int_{\Omega_t}(E) \cdot v dx \implies$

$$E_0(t) = \int_{\Omega_t} \frac{1}{2}|v|^2 dx + \alpha S(\partial\Omega_t) = E_0(0)$$

- $S(\partial\Omega_t)$: surface area \Rightarrow regularity control on $\partial\Omega_t$?
- Lagrangian formulation?

Lagrangian formulation

- Action:

$$\mathcal{A}(u) = \int \int_{\Omega_0} |u_t|^2 dy dt - \int \alpha S(\partial \Omega_t) dt$$

$u(t, \cdot) : \Omega_0 \rightarrow \Omega_t$: Lagrangian coordinate map, $u_t = v(t, u)$.

$u(t, \cdot) \in$ configuration space \mathcal{M} :

$$\mathcal{M} = \{u : \Omega_0 \rightarrow \mathbf{R}^n : \det \nabla u = 1\} \subset L^2(\Omega_0, \mathbf{R}^n)$$

$$T_u \mathcal{M} = \{W : \Omega_0 \rightarrow \mathbf{R}^n : \nabla \cdot (W \circ u^{-1}) = 0\}$$

$$(T_u \mathcal{M})^\perp = \{(\nabla q) \circ u : \Omega_0 \rightarrow \mathbf{R}^n : q|_{u(\partial \Omega)} = 0\}$$

- $S(\partial \Omega_t)$ depends only on $\partial \Omega_t = u(\partial \Omega_0)$

- Euler-Lagrangian equation:

$$u_{tt} + \alpha S'(u) = (\text{LM}) \triangleq \text{Lagrangian multiplier} \quad (\text{E})$$

- * $S'(u) \in T_u \mathcal{M}$: div free in Eulerian coordinates
- * $(\text{LM}) \in (T_u \mathcal{M})^\perp \Rightarrow (\text{LM}) = (\nabla p_0) \circ u, p_0|_{\partial\Omega} = 0$

$$\mathcal{D}_t u_t + \alpha S'(u) = 0, \quad (\text{E})$$

\mathcal{D} : covariant derivative on \mathcal{M}

Splitting of pressure

- $(LM) = -\nabla p_0$:

$$-\Delta p_0 = \partial_i v^j \partial_j v^i \quad p_0|_{\partial\Omega} = 0$$

* p_0 : volume preserving pressure

- $S' = -\nabla \tilde{p}$:

$$\Delta \tilde{p} = 0 \quad \tilde{p}|_{\partial\Omega_t} = \kappa : \text{mean curvature}$$

* \tilde{p} : surface tension

* $\forall w, \nabla \cdot w = 0$: $\langle S', w \rangle = \int_{\partial\Omega_t} \kappa w \cdot N dS = \int_{\Omega_t} \nabla \tilde{p} \cdot w dx$

- $p = p_0 + \alpha \tilde{p}$: reg. of $\tilde{p} <$ reg. of p_0

Linear analysis I.

- Linearized Euler-Lagrangian equation:

$$W_{tt} + \alpha S''(u)W = \dots$$

$W(t, \cdot)$: div free in Eulerian coordinates

- $S'' = A + l.o.t.$

* A self-adjoint on $L^2(\Omega_t)$ and $\forall w, \nabla \cdot w = 0$

$$\langle w, Aw \rangle_{L^2(\Omega_t)} = \int_{\Omega_t} (w, Aw) dx \triangleq \int_{\partial\Omega_t} |\nabla^{tan} w^\perp|^2 dS$$

* $w^\perp = w \cdot N$: normal velocity of $\partial\Omega$

Splitting of velocity

- For vector field w on Ω_t , $\nabla \cdot w = 0$,

$$w = w^{irrot} + w^{rot}$$

$$\nabla \cdot w^{rot} = 0 = \nabla \cdot w^{irrot}$$

- $w^{rot}|_{\partial\Omega_t} \in T\partial\Omega_t$: internal rotations
- $(w^{irrot})^\perp|_{\partial\Omega_t} = w^\perp|_{\partial\Omega_t}$: motion of $\partial\Omega_t$, $w^{irrot} = \nabla\phi$
- $\langle w^{rot}, w^{irrot} \rangle_{L^2(\Omega_t)} = 0$

Linear analysis II.

$$W_{tt} + \alpha S''(u)W = \dots$$

$W(t, \cdot)$: div free in Eulerian coordinates

- $S'' = A + l.o.t.$

* A self-adjoint on $L^2(\Omega_t)$ and $\forall w, \nabla \cdot w = 0$

$$\langle w, Aw \rangle_{L^2(\Omega_t)} = \int_{\partial\Omega_t} |\nabla^{tan} w^\perp|^2 dS \sim \|w^{irrot}\|_{H^{\frac{3}{2}}(\Omega_t)}^2$$

- $\partial_t \sim (\partial_x)^{\frac{3}{2}}$: only geometrically, may not be reflected in Lagrangian coordinates!

Energy estimates

$$u_{tt} + S'(u) = (\text{LM}) \quad (\text{E})$$

$A^k u_t \cdot (\text{E})$ & vorticity transport \Rightarrow

Theorem. (**Shatah-Z**) *Estimates (local in time) on*

$$v = u_t \circ u^{-1} \in H^{\frac{3}{2}k}(\Omega_t), \quad \partial\Omega_t \in H^{\frac{3}{2}k+1}, \quad p \in H^{\frac{3}{2}k-\frac{1}{2}}(\Omega_t)$$

- Estimate on $\partial\Omega_t$ in terms of κ , coordinate independent
- Interface problem (2F).

(1F): Without surface tension

$$v_t + v \cdot \nabla v = -\nabla p, \quad x \in \Omega_t \quad (\text{E})$$

$v|_{\partial\Omega_t}$: boundary velocity, $p|_{\partial\Omega_t} = 0$

- No $A = S'' + l.o.t.$ or \tilde{p}
- $p = p_0$: ∇p_0 becomes a leading order term

Geodesic Flow

Action: $\mathcal{A} = \int \int_{\Omega_0} |u_t|^2 dy dt, \ u(t, \cdot) \in \mathcal{M}$

$$\mathcal{M} = \{u : \Omega_0 \rightarrow \mathbf{R}^n : \det \nabla u = 1\} \subset L^2(\Omega_0, \mathbf{R}^n)$$

$$T_u \mathcal{M} = \{W : \Omega_0 \rightarrow \mathbf{R}^n : \nabla \cdot (W \circ u^{-1}) = 0\}$$

- $\frac{\delta \mathcal{A}}{\delta u} = 0$: geodesic flow on \mathcal{M}

$$\mathcal{D}_t u_t = 0,$$

\mathcal{D} : covariant derivative on \mathcal{M}

Linear Analysis: Jacobi Fields

- Linearized equation:

$$\mathcal{D}_t^2 W + \mathcal{R}(u_t, W)u_t = 0$$

\mathcal{R} : curvature of \mathcal{M} , for $W \in T_u \mathcal{M}$,

$$\langle \mathcal{R}(u_t, W)u_t, W \rangle = \int_{\partial\Omega_t} -\frac{\partial p}{\partial N} |W \cdot N|^2 dS + \dots \triangleq \langle \mathcal{R}_0(W), W \rangle + \dots$$

- $\mathcal{R}(u_t, \cdot)u_t$: unbounded \sim 1st order differential operator for W no smoother than $\nabla_N p$
- **[RT]** occurs unless $-\frac{\partial p}{\partial N} > 0$ on $\partial\Omega_t$: **(RT)** sign condition

Energy estimates

- $J = \nabla \tilde{p}$ is almost a Jacobi field (linearized solution):

$$\mathcal{D}_t^2 J + \mathcal{R}(u_t, J)u_t = l.o.t.$$

- $\mathcal{D}_t \kappa = \Delta_{\partial \Omega_t} v^\perp + l.o.t.$

(RT) Assume $-\frac{\partial p}{\partial N} > a > 0$ on $\partial \Omega_t$ for $t \in [0, T_0]$

Theorem. **(Shatah-Z)** Assume **(RT)**. Estimates on

$$v = u_t \circ u^{-1} \in H^k(\Omega_t), \quad \partial \Omega_t \in H^k, \quad p \in H^{k+\frac{1}{2}}(\Omega_t).$$

Uniform estimates hold as surface tension $\alpha \rightarrow 0$.

- **(2F)** w/o surface tension $\Rightarrow R(u_t, \cdot)u_t < 0$ 2nd order \Rightarrow **[KH]**.

- Surface tension vs. No surface tension (**1F**):

- * No surface tension:

$$\text{reg. of } \partial\Omega_t = \text{reg. of } v|_{\partial\Omega_t} + \frac{1}{2} \quad \text{reg. of } p = \text{reg. of } v + \frac{1}{2}$$

- * With surface tension:

$$\text{reg. of } \partial\Omega_t = \text{reg. of } v|_{\partial\Omega_t} + \frac{3}{2} \quad \text{reg. of } p = \text{reg. of } v - \frac{1}{2}$$

- Lagrangian coordinates optimal?

- * Example:

$$\Omega_t = B(1) \subset \mathbf{R}^2, \quad v(t, r, \theta) = \Theta(r) \frac{\partial}{\partial \theta}, \quad p(t, r, \theta) = \int_r^1 r' \Theta(r')^2 dr'$$

$$u(t, r_0, \theta_0) = (r_0, \theta_0 + t\Theta(r_0)), \quad \text{supp}(\Theta) \subset \subset (0, 1)$$

Construction of solutions (w. surface tension)

- Fix the moving domains with a nearby reference domain Ω_*^\pm
- * Represent $\partial\Omega_t$ by a modified mean curvature $\kappa_M : \partial\Omega_* \rightarrow \mathbf{R}$
- * Determine normal velocity (& thus v^{ir}) by $\partial_t \kappa_M$
- * Transfer v^{rot} to $v_*^{rot} : \Omega_* \rightarrow \mathbf{R}^n$ with $v_*^{rot}|_{\partial\Omega_*} \in T\partial\Omega_*$

1. Start with $(\kappa_M, \partial_t \kappa_M, v_*^{rot})$
2. Reconstruct (Ω_t, v) and u by solving elliptic equations
3. Solve for $(\tilde{\kappa}_M, \partial_t \tilde{\kappa}_M)$ from an equ derived from (L)

$$\partial_{tt} \tilde{\kappa}_M + "|\partial^3|"\tilde{\kappa}_M = Q(\kappa_M, \partial_t \kappa_M, v_*^{rot})$$
4. Construct \tilde{v}_*^{rot} by symmetries (Noether's Theorem)
5. contraction mapping

Theorem. **(Shatah-Z, 08)** (1F) is locally well-posed in the space $\partial\Omega \in H^{\frac{3}{2}k+1}$ and $v \in H^{\frac{3}{2}k}(\Omega_t)$.

- w/o surface tension assuming **(RT)**: vanishing surface tension limit
- **(2F)** w. surface tension

Other cases of (1F)

$$v_t + v \cdot \nabla v = -\nabla p - g e_n, \quad x \in \Omega_t \quad (\text{E})$$

* Ω_t : allowed to be infinite domain

* $\partial\Omega_t = S_t \cup B$, $d(S_t, B) \geq a > 0$

$$v \cdot N = 0 \text{ on } B, \quad p = \alpha \kappa \text{ on } S_t \quad (\text{BC})$$

* B : rigid part of boundary (bottom, obstacles ...)

* S_t : free surface, asymptotically flat in space

* g : gravity

Framework

- Action:

$$\mathcal{A}(u) = \int \int_{\mathbf{R}^n} \frac{1}{2} |u_t|^2 dy dt - \alpha \int S(u(S_0)) dt - G(u)$$

- * $u(t, \cdot)$: Lagrangian coordinate map

- * Gravitational potential:

$$G(u) = \int_{u(\Omega_0) \cap x_n > 0} x_n dx - \int_{u(\Omega_0) \cap x_n < 0} x_n dx$$

configuration space \mathcal{M} :

$$\mathcal{M} = \{u : \Omega_0 \rightarrow \mathbf{R}^n : \det \nabla u = 1, u(B) = B\} \subset L^2(\Omega_0, \mathbf{R}^n)$$

$$T_u \mathcal{M} = \{W : \Omega_0 \rightarrow \mathbf{R}^n : \nabla \cdot (W \circ u^{-1}) = 0, (W \cdot N)|_{u(S_0)} = 0\}$$

$$(T_u \mathcal{M})^\perp = \{(\nabla q) \circ u : \Omega_0 \rightarrow \mathbf{R}^n : q|_{u(\partial\Omega)} = 0\}$$

- Linearization:

$$\mathcal{D}_t^2 W + (\mathcal{R}(u_t, \cdot) u_t + D^2 G(u)) W + \alpha D^2 S(u) W = 0$$

- For $\alpha = 0$, **(RT)** from $\mathcal{R}(u_t, \cdot) u_t + D^2 G(u)$.

Framework for (2F)

- Action:

$$\mathcal{A}(u) = \int \int_{\mathbf{R}^n} \frac{\rho}{2} |u_t|^2 dy dt - \int S(u) dt$$

* $u(t, \cdot) = (u_+, u_-) : \mathbf{R}^n \setminus S_0 \rightarrow \mathbf{R}^n \setminus S_t$: Lagrangian coordinates

- Configuration space:

$$\begin{aligned} \mathcal{M} = \{u_\pm : \Omega_0^\pm \rightarrow \Omega_t^\pm : & \text{volume preserving diffeo. s.t.} \\ & S \triangleq u_+(S_0) = u_-(S_0)\} \end{aligned}$$

* In Eulerian coordinates:

$$T_u \mathcal{M} = \{v = (v_+, v_-) : \nabla \cdot v_\pm = 0, (v_+^\perp + v_-^\perp)|_S = 0\} \subset L^2(\rho dx)$$

$$(T_u \mathcal{M})^\perp = \{w = (\nabla \phi_+, \nabla \phi_-) : (\rho_+ \phi_+ - \rho_- \phi_-)_S = 0\} \subset L^2(\rho dx)$$