HIKARI

Site Search: [_____] Go

Keguang Liu, Xiaolong Qin
A gradient recovery-based a posteriori error estimators for the Ciarlet - Raviart formulation of the second biharmonic equations
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 997-1007.

H. Roumili, A. Keraghel, A. Yassine
Infeasible interior point method for semidefinite programs
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1009-1018.

M. Laghdir, R. Benabbou
Convex functions whose epigraphs are semi-closed: Duality theory
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1019-1033.

L. Bougoffa
A class of second order differential equations with nonlocal boundary conditions
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1035-1039.

L. Bougoffa
Parabolic equations with nonlocal conditions
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1041-1048.

L. Bougoffa
Modified decomposition method for solving the Cauchy problem for nonlinear parabolic-hyperbolic equations
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1049-1056.

J. A. Vera
Stability of a four body relative equilibrium
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1057-1067.

Weiming Ke
Optimal selection of blocked two-level fractional factorial designs
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1069-1082.

J. A. Diaz-Garcia, G. Gonzalez-Farias, V. M. Alvarado-Castro
Exact distributions for sensitivity analysis in linear regression
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1083-1100.

M. Laklalech, O. El Kahlaoui, M. Rachik, A. Namir, A. Bennar
On a class of integrodifferential equations
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1101-1120.

K. Al Shaqsi, M. Darus
On certain subclass of analytic univalent functions with negative coefficients
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1121-1128.

J. Penm
The recursive fitting of multivariate complex subset ARX models
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1129-1143.

Shenghui Chen, Qinghua Chen
A weighted network model based on the preferential selection of edges
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1145-1156.

D. M. Young, J. D. Stamey, J. W. Seaman, Jr.
Nonnegative-definite covariance structures for which the least squares estimator is the best linear unbiased estimator
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1157-1168.

M. B. Abd-el-Malek, N. A. Badran, H. S. Hassan

[Lie-group method for predicting water content for immiscible flow of two fluids in a porous medium](#)
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1169-1180.

N. Azizi, R. Pourgholi
[A nonlinear inverse parabolic problem](#)
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1181-1186.

[Lie-group method for predicting water content for immiscible flow of two fluids in a porous medium](#)
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1169-1180.

N. Azizi, R. Pourgholi
[A nonlinear inverse parabolic problem](#)
Appl. Math. Sci., Vol. 1, 2007, no. 21-24, 1181-1186.