

Picard group of hypersurfaces in toric varieties

Ugo Bruzzo, Antonella Grassi

(Submitted on 3 Nov 2010 (v1), last revised 12 Nov 2010 (this version, v2))

We show that the usual sufficient criterion for a generic hypersurface in a smooth projective manifold to have the same Picard number as the ambient variety can be generalized to hypersurfaces in complete simplicial toric varieties. This sufficient condition is always satisfied by generic K3 surfaces embedded in Fano toric 3-folds.

Comments: 14 pages. v2: some typos corrected

Subjects: **Algebraic Geometry (math.AG)**

MSC classes: 14C22, 14J70, 14M25

Report number: SISSA 78/2010/fm

Cite as: [arXiv:1011.1003v2](https://arxiv.org/abs/1011.1003v2) [math.AG]

Submission history

From: Ugo Bruzzo [[view email](#)]

[v1] Wed, 3 Nov 2010 20:23:53 GMT (12kb)

[v2] Fri, 12 Nov 2010 16:12:55 GMT (12kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AG

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1011](#)

Change to browse by:

[math](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

