

## Properties of Strange Matter in a Model with Effective Lagrangian

WANG Ping,<sup>1</sup> SU Ru-Keng,<sup>2,3</sup> SONG Hong-Qiu<sup>2,4,5</sup> and ZHANG Li-Liang<sup>4</sup>

<sup>1</sup> Institute of High Energy Physics, The Chinese Academy of Sciences, P.O. Box 918(4), Beijing 100039, China

<sup>2</sup> CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China

<sup>3</sup> Department of Physics, Fudan University, Shanghai 200433, China

<sup>4</sup> Shanghai Institute of Nuclear Research, The Chinese Academy of Sciences, P.O. Box 800204, Shanghai 201800, China

<sup>5</sup> Center of Nuclear Theory, National Laboratory of Heavy Ion Accelerator, The Chinese Academy of Sciences, Lanzhou 730000, China

(Received: 2000-4-30; Revised: )

**Abstract:** The strange hadronic matter with nucleons,  $\Lambda$ -hyperons and  $\Xi$ -hyperons is studied by using an effective nuclear model in a mean-field approximation. The density and strangeness fraction dependence of the effective baryon masses as well as the saturation properties and stabilities of the strange hadronic matter are discussed.

**PACS:** 21.65.+f, 21.80.+a, 21.10.Dr

**Key words:** strange hadronic matter, saturation, stability, effective Lagrangian

[\[Full text: PDF\]](#)

Close