



Mathematical Physics

An inequality for the trace of matrix products, using absolute values

Bernhard Baumgartner

(Submitted on 30 Jun 2011 ([v1](#)), last revised 1 Sep 2011 (this version, [v2](#)))

The absolute value of matrices is used in order to give inequalities for the trace of products. An application gives a very short proof of the tracial matrix Hoelder inequality

Comments: 5 pages

Subjects: **Mathematical Physics (math-ph)**

MSC classes: 39B42, 15A45, 47A50

Cite as: [arXiv:1106.6189](#) [math-ph]

(or [arXiv:1106.6189v2](#) [math-ph] for this version)

Submission history

From: Bernhard Baumgartner [[view email](#)]

[\[v1\]](#) Thu, 30 Jun 2011 11:21:43 GMT (5kb)

[\[v2\]](#) Thu, 1 Sep 2011 16:01:51 GMT (5kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math-ph

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[math](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

