



Control for Schroedinger operators on tori

Nicolas Burq, Maciej Zworski

(Submitted on 7 Jun 2011)

A well known result of Jaffard states that an arbitrary region on a torus controls, in the L2 sense, solutions of the free stationary and dynamical Schroedinger equations. In this note we show that the same result is valid in the presence of a smooth time-independent potential. The methods apply to continuous potentials as well and we conjecture that the L2 control is valid for any bounded time dependent potential.

Comments: 2 figures

Subjects: **Analysis of PDEs (math.AP)**; Systems and Control (cs.SY); Mathematical Physics (math-ph); Optimization and Control (math.OC)

MSC classes: 35Q41, 35Q93, 49J20, 81Q93

Cite as: **arXiv:1106.1412 [math.AP]**
(or **arXiv:1106.1412v1 [math.AP]** for this version)

Submission history

From: Maciej Zworski [[view email](#)]

[v1] Tue, 7 Jun 2011 18:41:21 GMT (20kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math.AP

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

- cs
- cs.SY
- math
- math-ph
- math.OC

References & Citations

- [NASA ADS](#)

Bookmark [\(what is this?\)](#)

