

Motif based hierarchical random graphs: structural properties and critical points of an Ising model*

Monika Kotorowicz[†], Yuri Kozitsky[‡]

Institute of Mathematics, Maria Curie-Skłodowska University, Lublin, Poland

Received March 15, 2010, in final form June 7, 2010

A class of random graphs is introduced and studied. The graphs are constructed in an algorithmic way from five motifs which were found in [Milo R., Shen-Orr S., Itzkovitz S., Kashtan N., Chklovskii D., Alon U., Science, 2002, **298**, 824–827]. The construction scheme resembles that used in [Hinczewski M., A. Nihat Berker, Phys. Rev. E, 2006, **73**, 066126], according to which the short-range bonds are non-random, whereas the long-range bonds appear independently with the same probability. A number of structural properties of the graphs have been described, among which there are degree distributions, clustering, amenability, small-world property. For one of the motifs, the critical point of the Ising model defined on the corresponding graph has been studied.

Key words: amenability, degree distribution, clustering, small-world graph, Ising model, critical point

PACS: 89.75.Fb, 89.75.Kd, 05.10.Cc, 05.70.Jk

1. Introduction and setup

A vast variety of large systems occurring in nature and society have a very complicated topological structure. These are the Internet, the World Wide Web, citation, neural, and social networks, etc. In view of the complex topology and unknown organizing principles, the networks are often modeled as random graphs. A random graph with a given node set V is a graph in which for a given pair $i, j \in V$, the bond $\langle i, j \rangle$ appears at random. The study of random graphs has been originated by P. Erdős and A. Rényi, who were the first to introduce such graphs in [1, 2]. In the Erdős-Rényi random graph model, denoted by $G_{n,p}$, the number of nodes is n , and the bonds between distinct nodes appear independently¹ with the same probability p . An important characteristic of a graph is the *node degree*, which is the number of bonds attached thereat. In $G_{n,p}$, it is a random variable, and all such variables are independent and have the same probability distribution. Namely, the probability that the degree of a given node is k is given by the Bernoulli law

$$P_n(k) = \binom{n-1}{k} p^k (1-p)^{n-1-k}, \quad k = 0, 1, \dots, n-1. \quad (1)$$

For random graphs, the most interesting questions refer to their asymptotic properties in the limit $n \rightarrow +\infty$. To get nontrivial answers to such questions one allows the parameters to depend on n . For $p_n = c/n$, the limit of (1) is the Poisson law

$$P(k) = c^k e^{-c} / k!, \quad k \in \mathbb{N}_0. \quad (2)$$

However, for irregular complex networks, the random graph $G_{n,p}$ is not a good model since in the most of such networks the degree distribution is essentially non-Poissonian. Many real world

*This work was supported by the DFG through the project 436 POL 125/0–1 as well as through SFB 701: “Spektrale Strukturen und Topologische Methoden in der Mathematik”. Yuri Kozitsky was also supported by TODEQ MTKD-CT-2005-030042.

[†]E-mail: monika@hektor.umcs.lublin.pl

[‡]E-mail: jkozi@hektor.umcs.lublin.pl

¹C.f., however, the discussion in [3].

networks, e.g. the WWW, are characterized by power-law node degree distributions, which have the form $P(k) = Ck^{-\gamma}$, $\gamma > 1$, typical for the so-called *scale-free* graphs.

Another important parameter characterizing a random graph is the *clustering coefficient*, which is the probability that two nodes are neighbors given they have a common neighbor. Clearly, in $G_{n,p}$ this probability is p , and is the same independently of whether or not the nodes have a common neighbor. Real world networks usually manifest strong clustering, which once more indicates that Erdős-Rényi type random graphs are not appropriate as their models. In [4], D.J. Watts and H. Strogatz proposed another type of random graphs, in which the mentioned disadvantage is overcome. It should be noted, however, that such graphs do not have power law node degree distributions. The next step beyond the Erdős-Rényi model was done by A.-L. Barabási and R. Albert in [5]. In their model, the preferential attachment principle has been employed, typical for many real networks (the Internet, citation and social networks). According to this principle, the more connected a node is, the more likely it receives a new bond. The construction of the Barabási-Albert model starts from an initial graph with $m_0 \geq 2$ nodes (neither can be isolated). At each step, one adds a new node and connects it to the existing nodes. The probability p_i that the new node is connected to node i is proportional to the degree n_i of that node, that is, $p_i = n_i / (\sum_j n_j)$.

Many of the complex networks occurring in nature contain characteristic patterns, recurring much more frequently than the other ones. They are called *network motifs*, see [6–11]. Different networks may have different motifs, and motifs in turn can characterize the networks. For instance, in biological regulation networks it has been experimentally demonstrated that each of the motifs can perform a key information processing function, see [8]. In [10], the authors introduced a random graph model based on some geometric principles (constraints). Then, they compared the appearance of eight elementary three- and four-node patterns in their model with the same characteristics of the Erdős-Rényi random graph. It turned out that five of these patterns are motifs for their model, but not for the Erdős-Rényi random graph, see figure 1 below and table 1 in [10]. For




Figure 1. Three and four node motifs M_1, M_2, M_3, M_4, M_5 found in [10].

another random graph model, the appearance of the same eight patterns was also studied in [11]. Note that among these patterns, only M_1 and M_5 correspond to complete graphs (each node is a neighbor to every other node).

One of the ways to get information about infinite graphs, also random ones, is to study the properties of certain models of statistical physics defined thereon. The most popular ones are the Ising and Potts models, as well as the models of bond and site percolation, see [12]. On the other hand, in statistical physics certain graphs are employed to mimic a crystal lattice, for which the critical-point behavior of the Ising model can be described in an explicit and rigorous way. These are the so-called *hierarchical lattices* introduced in [13]. Such lattices are defined in a rather algorithmic way by means of the basic pattern, e.g., by a “diamond”, which is the pattern M_3 depicted in figure 1. A mathematical description of the Gibbs states of the Ising model on such graphs was done by P.M. Bleher and E. Žalys in [14, 15]. M. Hinczewski and A. Nihat Berker [16] studied the critical-point properties of the Ising model on the diamond hierarchical lattice ‘decorated’ by additional bonds, which appear at random. In the present paper, we follow the way suggested in [16] and introduce hierarchical graphs constructed by means of the motifs shown in figure 1, decorated by additional bonds which appear at random and repeat, in a way, the corresponding motif. We analyse some of their characteristics, such as the average degree, the node degree distribution, amenability, the small-world property, as well as the critical-point properties of the Ising model. This study of ours is a continuation of [17], where the graph based on M_1 was introduced by one of us. Note that our hierarchical graphs also found applications in cryptography, see [18].

2. The graphs: construction and structural properties

2.1. The construction: informal description

As is typical for hierarchical graphs, e.g., for hierarchical lattices in [13, 16], the construction is carried out in an algorithmic way: at k -th level, $k \in \mathbb{N}$, one produces a subgraph, say Λ_k , which is then used as a construction element for producing Λ_{k+1} . The procedure is the same at each level, and the starting element is obtained from the corresponding motif. Let us illustrate this for the simplest case based on the motif M_1 . First we label the nodes of M_1 by a , b , and c , as it is shown in figure 2, and obtain Λ_1 — the triangle. Then we take three such graphs and label them by the same labels. As a result, each of the triangles has nodes of two kinds: one node the label of which coincides with the triangle label (e.g., node a in triangle a), and two nodes with non-coinciding labels. Thereafter, the triangles are glued up according to the following rule: node c of triangle a is glued up with node a of triangle c , ect. The nodes with the coinciding labels remain untouched. These are the so-called *external* nodes of Λ_2 . The remaining nodes are called *internal*. The bonds of the initial triangles turn into the bonds of Λ_2 . We call them *basic* bonds; they are depicted by solid lines. At the next stage of the first step, we add the bonds connecting the external nodes in the same way as it is in the motif M_1 . Such bonds are depicted by dotted lines and are called *decorations*. As a result, we obtain the graph Λ_2 , which has six basic bonds and three decorations, three external and three internal nodes. Then we repeat the same procedure — take three copies of Λ_2 , label them by a , b , and c , and divide their external nine nodes into two groups: three nodes with coinciding labels and six nodes with non-coinciding labels. Then the graphs Λ_2 are glued up as described above. Thereafter, three decorating bonds are drawn to connect the external nodes. This procedure is repeated ad infinitum. Similar constructions for motifs M_2 and M_3 are presented in figure 3 and figure 4, respectively. Here we have omitted the decorating bonds not to overload the pictures. The picture for M_4 is obtained from that for M_3 by adding the diagonals. The picture for M_5 is just the three-dimensional version of the picture for M_1 , where the basic pattern is a tetrahedron.

2.2. Definitions

In order to fix the terminology and to make the construction of our graphs mathematically immaculate, we begin by introducing a number of relevant mathematical notions. A (simple) graph G is a pair of sets (V, E) , where V is the set of nodes, whereas E is a subset of the Cartesian product $V \times V$. We suppose that E is symmetric and irreflexive, i.e., $\langle j, i \rangle \in E$ whenever $\langle i, j \rangle \in E$, and $\langle i, i \rangle \notin E$ for every $i, j \in V$. We say that i and j are connected by a *bond* if $\langle i, j \rangle \in E$. In this case, we write $i \sim j$ and say that i and j are *adjacent* or that they are *neighbors*. Hence, the elements of E themselves can be called bonds. The graph is said to be *complete* if each two nodes are adjacent. For a given i , by $n(i)$ we denote the *degree* of i — the number of its neighbors. If V is finite, then E is also finite, and the graph is said to be finite. Otherwise, the graph is infinite. An infinite graph is called *locally finite*, if $n(i)$ is finite for every node. All the infinite graphs we study are *countable*, which means that both sets V and E are infinite and countable. Given $G = (V, E)$ and $G' = (V', E')$, let $\phi : V \rightarrow V'$ be such that $\phi(i) \sim \phi(j)$ whenever $i \sim j$. Such a map ϕ is called a *morphism*. A bijective morphism is called an *isomorphism*. If ϕ is an isomorphism, then its inverse ϕ^{-1} is also an isomorphism, and then the graphs G and G' are said to be mutually isomorphic. Such graphs have identical structures and thus can be identified. In this case, we also say that G' is a *copy* of G . One observes that this refers to both finite and infinite graphs. An isomorphism $\phi : V \rightarrow V$, i.e. which maps the graph onto itself, is called an *automorphism*. The graph $G' = (V', E')$ such that $V' \subset V$ and $E' \subset E$ is said to be a *subgraph* of $G = (V, E)$. In this case, we write $G' \subset G$. Suppose that a subgraph $G' \subset G$ has a copy, say \tilde{G} , that is, there exists an isomorphism $\phi : \tilde{G} \rightarrow G'$. Then ϕ , considered as a map $\phi : \tilde{G} \rightarrow G$, is called an *embedding* of \tilde{G} into G , whereas G' is called the *image* of \tilde{G} under this embedding. Figure 1 presents the so-called *unlabeled* graphs, which we call patterns. After labeling, i.e., attaching a label to each of the nodes, such a pattern turns into a graph. Another labeling may or may not yield the same graph. This depends on whether or not

there exists the corresponding automorphism. For instance, any labeling of the triangle M_1 yields the same graph since in any case each of the nodes has the same neighbors. For the pattern M_2 , the left-hand graph in figure 3 with the interchanged labels a and b is the same. However, the graph with the interchanges c and d is not the same anymore. Of course, this new graph is isomorphic to the initial one. This is because there is only one nontrivial automorphism of M_2 : the one which interchanges a and b , and preserves c and d . The triangle has six automorphisms.

Let us now turn to *random* graphs. To introduce such a graph we need an *underlying* graph $G = (V, E)$ and a family \mathcal{E} of subsets of E . If G is finite, as \mathcal{E} one can take the set of all subsets of E . In the sequel, we deal with such graphs only. Thus, for $E' \in \mathcal{E}$, we say that E' has been picked *at random* with probability $P(E')$. In the Erdős-Rényi model $G_{n,p}$, the underlying graph is complete with the node set $V = \{1, \dots, n\}$. In this model, $P(E') = p^{|E'|}$, where $p \in [0, 1]$ and $|E'|$ stands for the number of elements in E' . In other words, the elements of E are being picked independently, each with the same probability p . In a bit complicated model, the bonds are picked independently but with probability which depends on the bond. In this case, as well as in the case of the Erdős-Rényi model, we deal with a random graph with independent bonds. For such graphs,

$$P(E') = \prod_{e \in E'} p(e), \quad (3)$$

where $p(e)$ is the probability of picking bond e . The set of graphs $G' = (V, E')$ with $E' \in \mathcal{E}$ is called the graph *ensemble* — each G' is picked at random from this ensemble. A *random graph model* is the pair consisting of the graph ensemble and of the function $P : \mathcal{E} \rightarrow [0, 1]$. If the function P is as in (3), the graph is said to be a random graph with independent bonds. Suppose that we have two random graph models with independent bonds. Let also $G = (V, E)$ and $\tilde{G} = (\tilde{V}, \tilde{E})$ be their underlying graphs and $\phi : V \rightarrow \tilde{V}$ be a morphism. Then this map is said to be the *morphism of the random graphs* if for every $\langle i, j \rangle \in E$, the probability (in the first model) that this bond is picked is the same as the corresponding probability (in the second model) for the bond $\langle \phi(i), \phi(j) \rangle$.

2.3. The construction

As was mentioned above, each of our graphs is constructed in an algorithmic way from the corresponding motif presented in figure 1. Since they are going to be random graphs with independent bonds, we have to construct the corresponding underlying graphs and, to define, for a given bond, the probability of being picked, c.f. (3). In all our models, the bonds will be of two kinds, which we call basic bonds and decorations. Basic bonds are going to be non-random, i.e. they are picked with probability one. Decorating bonds appear with probability $p \in [0, 1]$, which is a parameter of the model. Turn now to the construction of the underlying graphs. Let q be the number of nodes in the corresponding motif, that is, $q = 3$ for M_1 and $q = 4$ for the remaining motifs. At step $k = 1$, we just label the vertices of the corresponding motif by $i = 1, \dots, q$ and obtain the initial graph $\Lambda_1 = (V_1, E_1)$. All its bonds are set to be basic. Suppose now that we have $q + 1$ copies of Λ_1 obtained by the isomorphisms ϕ_2^j , $j = 0, 1, \dots, q$. Thus, in j -th copy the nodes are $\phi_2^j(i)$, $i = 1, \dots, q$. The graph Λ_2 is obtained from these copies under the following conditions

$$\phi_2^0(i) = \phi_2^i(i), \quad i = 1, \dots, q; \quad \phi_2^i(j) = \phi_2^j(i), \quad i = 1, \dots, q, \quad i \neq j. \quad (4)$$

Thus, the images of V_1 under ϕ_2^i and ϕ_2^j with $i \neq j$ intersect only at one node where (4) holds. The maps ϕ_2^j , $j = 0, 1, \dots, q$ embed Λ_1 into Λ_2 . The nodes $\phi_2^i(i)$, $i = 1, \dots, q$, are called the *external* nodes of Λ_2 . All other nodes are called *internal*. Thus, Λ_2 has q external and $q(q - 1)/2$ internal nodes. At this stage, we label them by $i = 1, \dots, q(q + 1)/2$ in such a way that the external nodes have the same labels as in Λ_1 , that is, $\phi_2^i(i) = i$, $i = 1, \dots, q$. By construction, the bonds obtained as images under the map ϕ_2^0 are decorations: they are of the form $\langle \phi_2^0(i), \phi_2^0(j) \rangle$ where i and j are adjacent in Λ_1 . From the first condition in (4) we see that the decorating bonds connect the external nodes of Λ_2 . The remaining bonds of Λ_2 are set to be basic. Now we construct Λ_3 from one copy of Λ_1 and q copies of Λ_2 . Let ϕ_3^0 be the map which produces the copy of Λ_1 and ϕ_3^j ,

$j = 1, \dots, q$ be the maps which produce the copies of Λ_2 . We then impose the conditions

$$\phi_3^0(i) = \phi_3^i(i), \quad i = 1, \dots, q; \quad \phi_3^i(j) = \phi_3^j(i), \quad i = 1, \dots, q, \quad i \neq j \quad (5)$$

and obtain Λ_3 . Thus, ϕ_3^0 embeds $\Lambda_1 \rightarrow \Lambda_3$, and $\phi_3^i : \Lambda_2 \rightarrow \Lambda_3$, $i = 1, 2, \dots, q$. As above, the nodes $\phi_3^i(i)$ are set to be external, and the remaining nodes are internal. The images of V_2 under ϕ_3^i and ϕ_3^j with $i \neq j$ intersect only at one node where (5) holds. Again we label the nodes of Λ_3 in such a way that $\phi_3^i(i) = i$, $i = 1, \dots, q$. Now let us establish which bonds of Λ_2 are decorating and which are basic. As above, the bonds connecting the external nodes are decorating. The images of decorating bonds of Λ_2 are decorating bonds in Λ_3 ; the same is also true for the basic bonds — the basic bonds of Λ_3 are exactly the images of the basic bonds of Λ_2 . For $k \geq 4$, the construction of Λ_k from Λ_{k-1} and Λ_1 is identical to the construction of Λ_3 just described. As above, by V_k and E_k we denote the sets of nodes and bonds of Λ_k , respectively. Thus, for $k \geq 2$ we have $E_k = E'_k \cup E''_k$, where E'_k (respectively, E''_k) consists of basic (respectively, decorating) bonds. All Λ_k , $k \in \mathbb{N}$, are considered as subgraphs of an infinite graph Λ_∞ , the structure and properties of which are not important for the study presented in this article.

Note that the construction principle used above essentially differs from that used in the construction of the hierarchical lattices in [13–16]. Namely, in our case to obtain Λ_k one replaces each *node* of the basic pattern by a copy of the graph Λ_{k-1} . In the hierarchical lattices, one replaces a *bond*. As we shall see in the sequel, this leads to essentially different properties of the resulting graphs. Below in figure 2, we illustrate the construction described above for the case where the basic pattern is the motif M_1 . In this case, the bare graph (i.e. the one which occurs for $p = 0$) is the approximating graph for the fractal known as the Sierpiński gasket². The elements of E'_2 (middle graph) and of E'_3 (right-hand graph) are depicted by solid lines. The elements of E''_2 and of E''_3 are depicted by dotted lines. We omit some dotted lines to indicate that they appear at random and hence may be absent in a given realization of the graph. Note that Λ_3 can be viewed as the triangle composed of three copies of Λ_2 . In figure 3, we present the construction of the bare graph Λ_3 corresponding to M_2 . In contrast to the former case, this is not a planar graph. In figure 4, we construct the bare graph Λ_2 for the motif M_3 . One observes that in that picture the node c of the lower left-hand quadrat (i.e. quadrat a) is glued up with node a of the upper right-hand quadrat. It is interesting that the corresponding fractal can be obtained by the following procedure, resembling the one which yields the Sierpiński gasket. One takes the full quadrat and cuts it out into four equal quadrats, but without cutting the external lines. Then, one glues up the vertices of the smaller quadrats as depicted and proceeds with cutting out the smaller quadrats. The fractal which one obtains from M_5 is a three dimensional version of the Sierpiński gasket. One takes the full tetrahedron and cuts out its inner one fourth in such a way that the remaining four tetrahedra are glued up according to the rule: vertex b of tetrahedron a is glued up with vertex a of tetrahedron b , etc.




Figure 2. Construction of the graph Λ_3 based on M_1 .

²The fractal itself is obtained as the closure of the set $\cup_{k \in \mathbb{N}} V_k$ in the appropriate topology, see e.g. [19].




Figure 3. Construction of the bare graph Λ_3 based on M_2 .




Figure 4. Construction of the bare graph Λ_2 based on M_3 .

2.4. Degree distribution

Now we turn to the description of the structural properties of the graphs constructed above. Let $m_k, k \in \mathbb{N}$, be the number of times the basic pattern appears in non-decorated Λ_k as a subgraph. For the graphs based on M_1 and M_5 we have the same situation. Here $m_1 = 1$ and $m_k = qm_{k-1}$, for $k \geq 2$ with the exception in Λ_2 , where the additional pattern appears. So, for M_1 and M_5 we obtain $m_1 = 1$ and

$$m_k = (q + 1)q^{k-2}, \quad k \geq 2. \quad (6)$$

Here $q = 3$ and $q = 4$ for M_1 and M_5 , respectively. For motif M_3 we have $m_1 = 1, m_2 = 2q$ and

$$m_k = qm_{k-1}, \quad k \geq 3. \quad (7)$$

Hence $m_k = 2 \cdot 4^{k-1}$ for $k \geq 2$. The simplest case is for M_4 , where $m_k = qm_{k-1}$ for $k \geq 2$. It gives $m_4 = 4^{k-1}$. The last motif is M_2 – triangle with additional bond. On each level this bond “produces” new 17 patterns. So $m_1 = 1$ and for $k \geq 2$ $m_k = \frac{1}{3}(26 \cdot 4^{k-1} - 17)$.

Now we analyze the number of times the basic pattern appears in fully decorated Λ_k , denoted by \tilde{m}_k . For the graphs based on M_1 and M_5 we have

$$\tilde{m}_k = \frac{2q + 1}{q - 1}q^{k-1} + \frac{q + 2}{q - 1}. \quad (8)$$

For M_3 it is $\frac{2}{3}4^k - \frac{5}{3}$ and for M_4 we obtain $\frac{1}{3}(4^k - 1)$. In all cases, we have m_k increasing as $C(p)q^{k-1}$, which means that adding decorations does not change the asymptotics of m_k .

In a similar way, we obtain

$$|V_k| = \frac{q^k + q}{2}, \quad |E_k| = rq^{k-1} + rp \frac{q^{k-1} - 1}{q - 1}, \quad k \in \mathbb{N}, \quad (9)$$

where $|V_k|$ stands for the number of nodes in Λ_k , whereas $|E_k|$ is the expected number of bonds in this graph.

As was mentioned above, the degree distribution is a very important characteristic of the graph. In contrast to Erdős-Rényi type graphs, for our graphs the distribution of the random variable $n(i)$ depends on the type of i . Thus, the simplest way to describe this distribution is to average $n(i)$ over the nodes of a given Λ_k , that is, to consider

$$n_k = \frac{1}{|V_k|} \sum_{i \in V_k} n(i). \quad (10)$$

Let $\langle n_k \rangle$ be the expected value of n_k . Then

$$\langle n_k \rangle = 2|E_k|/|V_k| = \frac{4r}{q(q-1)}(q-1+p) - \frac{4r}{q(q-1)} \cdot \frac{q-1+2p}{q^{k-1}+1}. \quad (11)$$

However, this result provides only partial information on the node degree distribution. To get more, let us analyze the structure of the node sets V_k , $k = 1, 2, \dots$, more in detail. For a given Λ_k and $l = 1, \dots, k$, let $V_k^{(l)}$ be the set of nodes $i \in V_k$ which are external for some Λ_l and, at the same time, are internal for any Λ_{l+1} . Of course, here we mean those Λ_l 's which are subgraphs for Λ_k . As an example, let us consider the graph Λ_2 based on M_1 , see the middle graph in figure 2. The nodes a , b , and c constitute $V_2^{(2)}$, whereas the remaining nodes constitute $V_2^{(1)}$.

The elements of $V_k^{(k-1)}$ are exactly the nodes at which the subgraphs Λ_{k-1}^j , $j = 1, \dots, q$ are glued up to form Λ_k , whereas the elements of $V_k^{(k)}$ are exactly the external nodes of Λ_k . Then $|V_k^{(k)}| = q$ and $|V_k^{(k-1)}| = q(q-1)/2$. For $l < k-1$, we have $|V_k^{(l)}| = q|V_{k-1}^{(l)}|$, which can be solved to yield

$$|V_k^{(l)}| = \frac{1}{2}q^{k-l}(q-1), \quad l = 1, \dots, k-1, \quad |V_k^{(k)}| = q. \quad (12)$$

The reason to consider the sets $V_k^{(l)}$ is that all the elements of each such $V_k^{(l)}$ have the same degree distribution, independent of k for $l \leq k-1$. In fact, the degrees of $i \in V_k^{(1)}$ are non-random since these nodes receive no decorating bonds. For such i , $n(i) = \sum_j n^{(0)}(j)$, where $n^{(0)}(j)$ is the degree of the corresponding node in the basic pattern, and the sum is taken over all such patterns which are glued up. By the symmetry of M_1 , M_3 , and M_5 , we have $n(i) = 4$ for M_1 and M_3 , and $n(i) = 6$ for M_5 . For M_2 , $n(i)$ takes values 3, 4, 5, see figure 3. For $i \in V_k^{(l)}$, $l = 2, 3, \dots, k-1$, we have $n(i) = \tilde{n}(i) + \nu(i)$, where $\tilde{n}(i)$ is non-random and has to be calculated as just described. The summand $\nu(i)$ is the number of decorating bonds attached to i . To simplify our consideration, let us stick to the case of the graph generated by M_1 . Then for $l = 1, \dots, k-1$ and $i \in V_k^{(l)}$, we have $\tilde{n}(i) = 4$ and $\nu(i)$ takes values $\nu = 0, 1, 2, \dots, 4(l-1)$, with probability

$$\text{Prob}(\nu(i) = \nu) = \binom{4(l-1)}{\nu} p^\nu (1-p)^{4(l-1)-\nu}. \quad (13)$$

For $i \in V_k^{(k)}$, $\nu(i)$ takes values $0, 1, \dots, 2(k-1)$. Therefore, the maximum node degree which can occur in V_k , $k \geq 2$, is

$$\max_{i \in V_k} n(i) = 4k - 4. \quad (14)$$

As is usual in the theory of real world networks, which are in fact non-random, the randomness manifests itself as the random choice of a node. If we apply this principle here, then (13) can be considered as the conditional probability distribution, conditioned by the event that the node i

has been picked from the set $V_k^{(l)}$. The probability of the latter event is taken to be proportional to the number of its elements, that is,

$$\text{Prob}\left(i \in V_k^{(l)}\right) = \frac{|V_k^{(l)}|}{|V_k|} = \frac{q-1}{1+q^{1-k}}q^{-l} = \frac{2}{1+3^{1-k}}3^{-l}, \quad l \leq k-1, \quad (15)$$

$$\text{Prob}\left(i \in V_k^{(k)}\right) = \frac{2q}{q^k+q} = \frac{2}{3^{k-1}+1}.$$

If we now take the expectation of $n(i)$ with respect to this distribution, that is,

$$\begin{aligned} \langle n_k \rangle &= \sum_{l=1}^{k-1} \sum_{\nu=0}^{4(l-1)} (4+\nu) \frac{2 \cdot 3^{-l}}{1+3^{1-k}} \cdot \binom{4(l-1)}{\nu} p^\nu (1-p)^{4(l-1)-\nu} \\ &+ \sum_{\nu=0}^{2(k-1)} (2+\nu) \frac{2}{3^{k-1}+1} \cdot \binom{2(k-1)}{\nu} p^\nu (1-p)^{2(k-1)-\nu}, \end{aligned} \quad (16)$$

we readily obtain $\langle n_k \rangle = 4 + 2p - 4(1+p)/(3^{k-1} + 1)$, which is in full agreement with the one given in (11) or in table 1. In order to figure out the limit $k \rightarrow +\infty$ of the distribution given by (13) and (15) we calculate its characteristic function, c.f. (16),

$$\begin{aligned} \varphi_k(t) &= \sum_{l=1}^{k-1} \sum_{\nu=0}^{4(l-1)} \exp[it(4+\nu)] \frac{2 \cdot 3^{-l}}{1+3^{1-k}} \cdot \binom{4(l-1)}{\nu} p^\nu (1-p)^{4(l-1)-\nu} \\ &+ \sum_{\nu=0}^{2(k-1)} \exp[it(2+\nu)] \frac{2}{3^{k-1}+1} \cdot \binom{2(k-1)}{\nu} p^\nu (1-p)^{2(k-1)-\nu} \\ &= \frac{2e^{4it}}{1+3^{1-k}} \cdot \frac{1-3^{1-k}(e^{it}p+1-p)^4}{3-(e^{it}p+1-p)^4} \\ &+ \frac{2e^{2it}}{3^{k-1}+1} (e^{it}p+1-p)^2, \quad i = \sqrt{-1}. \end{aligned} \quad (17)$$

Then the limiting characteristic function is

$$\varphi(t) = \frac{2e^{4it}}{3-(e^{it}p+1-p)^4}, \quad (18)$$

which can be continued to a meromorphic function analytic in some complex neighborhood of the point $t = 0$. This means that the limiting node degree distribution has all moments and hence cannot be of scale-free type³. This also agrees with the heuristic rule, see equation (3.13) on page 188 in [20], that for scale-free networks

$$\max_{i \in V_k} n(i) \sim |V_k|^{1/(\alpha-1)}, \quad \alpha > 1, \quad (19)$$

whereas in our case we have (14) and $|V_k| = (3^k + 3)/2$. In a similar way, one can show that all our graphs are not scale-free. Another observation on this item can be made by comparing the function (18) with the characteristic function of the Poisson distribution (2) which has the form

$$\varphi_{\text{Poisson}}(t) = \exp[c(e^{it} - 1)],$$

and hence can be continued to a function analytic on the whole complex plane. Therefore, the distribution corresponding to (18) with $p > 0$ is intermediate as compared to the Poisson and scale-free distributions. For $p = 0$, the function (18) is also entire.

³For scale-free graphs, the node degree distribution is $P(k) = Ck^{-\gamma}$, $k \geq 1$, $\gamma > 1$; hence, $\sum_{k=1}^{\infty} k^m P(k)$ diverges for all $m \geq \gamma - 1$.

2.5. Amenability, clustering, and small world properties

The next property of our graphs which we are going to address is *amenability*. To introduce it we need one more notion. Let $G = (V, E)$ be a countable graph with node set V and bond set E . For a finite $\Delta \subset V$, by $\partial\Delta$ we denote the set of nodes which are not in Δ but have neighbors in Δ . This set is the *outer boundary* of Δ , whereas the elements of Δ which are neighbors to the elements of $\partial\Delta$ constitute the *inner boundary* of Δ . As usual, by $|\Delta|$ and $|\partial\Delta|$ we denote the number of nodes in these sets. The graph G is said to be *amenable* if there exists a sequence of finite node sets $\{\Delta_k\}_{k \in \mathbb{N}}$, such that

$$\lim_{k \rightarrow +\infty} \frac{|\partial\Delta_k|}{|\Delta_k|} = 0. \quad (20)$$

If such a limit is positive for any sequence $\{\Delta_k\}_{k \in \mathbb{N}}$, the graph is called *nonamenable*. The lattices \mathbb{Z}^d , $d \geq 1$, are amenable graphs and for such sets one can take the cubes

$$\Delta_k = \{i = (i_1, \dots, i_d) \in \mathbb{Z}^d : |i_j| \leq N_k, j = 1, \dots, d\},$$

such that $N_{k+1} > N_k$ and $N_k \rightarrow +\infty$. In this case, $|\Delta_k| \sim N_k^d$ and $|\partial\Delta_k| \sim N_k^{d-1}$ and hence (20) holds. Sometimes, sequences for which (20) holds are called *Van Howe* sequences, see e.g. [21]. Cayley trees, except for \mathbb{Z} , are nonamenable. Let us turn now to our graphs. Due to their hierarchical structure, it is convenient to check (20) for the sequence of node sets of Λ_k , that is for $\{V_k\}_{k \in \mathbb{N}}$. By the construction of Λ_k , the inner boundary of each V_k is exactly the set of all its external nodes, the number of which is equal to the number of nodes in the corresponding motif, i.e. it is q . By construction, one of them becomes an external node of Λ_{k+1} , and the remaining $q - 1$ ones become inner nodes of Λ_{k+1} . For all the motifs M_j , $j = 1, \dots, 5$, we have the degree of any node therein being at most 3, see figure 1. Then for all our graphs,

$$\max_{i \in V_k} n(i) \leq 6k,$$

c.f. (14). At the same time, $|V_k| \sim q^k$, $q = 3, 4$, see (9), which immediately yields that all our graphs are amenable.

Now we study the clustering in our graphs. For non-random graphs, the clustering coefficient is defined as follows. For a given node $i \in V$ of degree $n(i)$, let $N(i)$ be the number of bonds linking its neighbors, which is the number of triangles with vertex i . Clearly, $N(i) \leq n(i)[n(i) - 1]/2$ and the maximum value of this parameter is attained for complete graphs where each node is a neighbor to any other one. Thus, the quantity

$$Q(i) := \frac{2N(i)}{n(i)[n(i) - 1]}, \quad (21)$$

characterizes clustering at node i . Then we define the clustering of our graphs as

$$Q = \lim_{k \rightarrow +\infty} \frac{1}{|V_k|} \sum_{i \in V_k} Q(i). \quad (22)$$

Note that for many graphs, e.g., for trees or bipartite graphs, one has $Q(i) = 0$ for any node i , see also [22, 23]. For random graphs, the degree $n(i)$, as well as the parameter $N(i)$, are random. Since in this case the calculation of Q is much more involved, we postpone it to the future. Here we only compare the values of Q obtained for the bare graphs with those for fully decorated ones.

For the graph based on M_1 and a node $i \in V_k^{(l)}$, $l = 1, \dots, k-1$, we have: (a) $n(i) = 4$, $N(i) = 3$ for $l = 1$, and $N(i) = 2$ for $l \geq 2$; (b) $n(i) = 4l$, $N(i) = 4l$. Here (a) and (b) correspond to a bare graph and to a fully decorated graph, respectively. These numbers follow directly from the construction of the graphs. The number of elements in each $V_k^{(l)}$ is given in (12), which allows one

to compute

$$(a) \quad Q = \lim_{k \rightarrow +\infty} \left(\frac{1}{3} + \frac{|V_k^{(1)}|}{6|V_k|} + \frac{2}{|V_k|} \right) = \frac{4}{9} = 0.4444\dots, \quad (23)$$

$$(b) \quad Q = 2 \cdot 3^{-1/4} \arctan 3^{-1/4} - \frac{1}{3^{1/4}} \ln \frac{3^{1/4} + 1}{3^{1/4} - 1} \approx 0.525897. \quad (24)$$

For the bare graph based on M_3 , we have $N(i) = 0$ for all nodes; hence, $Q = 0$ in this case. For the fully decorated graph based on M_3 , we observe that two nodes of $V_2^{(1)}$ have neighbors only in $V_2^{(1)}$, and the remaining four nodes have two neighbors in $V_2^{(1)}$ and two — in $V_2^{(2)}$. Since for the nodes $i \in V_2^{(1)}$, we have $N(i) = 0$, like in the case of the bare graph, we should consider these two groups separately. Thus, we split each $V_k^{(l)}$, $l = 1, \dots, k-1$, into $V_{k,0}^{(l)}$ and $V_{k,1}^{(l)}$, where the first set consists of the nodes which have neighbors only in $V_k^{(l)}$. The elements of $V_{k,1}^{(l)}$ have also neighbors in $V_k^{(l')}$ with $l' > l$. The number of nodes in these sets are:

$$|V_{k,0}^{(l)}| = \frac{1}{2} 4^{k-l}, \quad |V_{k,1}^{(l)}| = 4^{k-l}.$$

For all $i \in V_k^{(l)}$, $l = 1, \dots, k-1$, we have $n(i) = 4l$. At the same time, $N(i) = 4(l-1)$ for $i \in V_{k,0}^{(l)}$, and $N(i) = 1 + 4(l-1)$ for $i \in V_{k,1}^{(l)}$. Putting all these numbers together we obtain

$$Q = \frac{3}{2} \sum_{l=2}^{\infty} \frac{(l-1)4^{-(l-1)}}{l(4l-1)} + \sum_{l=1}^{\infty} \frac{4^{-l}}{l(4l-1)} \approx 0.1223. \quad (25)$$

For the bare graph based on M_5 one can obtain for $i \in V_k^{(l)}$, $l = 1, 2, \dots, k-1$: $n(i) = 6$, $N(i) = 8$ for $l = 1$, and $n(i) = 6$, $N(i) = 6$ for $l \geq 2$. For the fully decorated graph based on M_5 we have for all internal nodes $n(i) = 6l$, and $N(i) = 9$ for $l = 1$, and $N(i) = 12l + 7$ for $l \geq 2$. Hence, we obtain

$$(a) \quad Q = \lim_{k \rightarrow +\infty} \left(\frac{2}{5} + \frac{2|V_k^{(1)}|}{15|V_k|} + \frac{12}{5|V_k|} \right) = 0.5, \quad (26)$$

$$(b) \quad Q \approx 0.554145.$$

There exists one more property of real world networks, which Erdős-Rényi type graph does not share, see e.g. [20, 24]. It is the so-called *small-world property*, illustrated by Pal Erdős himself in the following way. All authors of mathematical papers are given the *Erdős index* which for his coauthors is equal to one. For their coauthors who are not Erdős' coauthors, this index is equal to 2, and so on. Everyone whose papers are indexed by the Mathematical Reviews can check his own index at the web-site <http://www.ams.org/mathscinet/>. It turns out that for many authors it ranges from 2 to 6 (for the second named author of this paper it is 4). To formulate the small-world property one needs the following notion. A path in the graph is a sequence of nodes such that every two consecutive elements of this sequence are neighbors to each other. The length of the path is the number of such consecutive pairs, which is equal to the number of bonds one passes on the way from the origin to the terminus. If every two nodes can be connected by a path, the graph is said to be connected. For the given two nodes, i and j , the length of the shortest path which connects them is said to be the distance $\rho(i, j)$ between these nodes. Informally speaking, a graph $G = (V, E)$ has the small-world property (or is a small-world graph) if every two nodes $i, j \in V$ are "not too far" from each other. More precisely this property is formulated as follows. An infinite graph G has a small-world property if there exists a sequence of its connected finite subgraphs $\{G_k\}_{k \in \mathbb{N}}$ with the following property. Let $\text{diam}(G_k) = \max_{i, j \in V_k} \rho(i, j)$ be the *diameter* of G_k , $k \in \mathbb{N}$, and $\langle n_k \rangle$ be the average value of the node degree in G_k , that is, $\langle n_k \rangle = 2|E_k|/|V_k|$. Then

the sequence $\{G_k\}_{k \in \mathbb{N}}$, and hence the graph G , are said to have the small-world property if there exists a positive constant C such that for all $k \in \mathbb{N}$,

$$\text{diam}(G_k) \leq C \log_{\langle n_k \rangle} |V_k|. \quad (27)$$

This means that in such graphs, the distances between the nodes scale at most logarithmically with the size of the graph. For our graphs, the results on this item are given in the last three rows of table 1. The diameters of Λ_k presented therein were calculated in a routine way for the cases where the graphs are bare ($p = 0$) or are fully decorated ($p = 1$). In the former case, neither of our graphs has the small-world property. However, this property holds for all fully decorated graphs.

Table 1. The structural characteristics of the families of hierarchical graphs.

motif	M_1	M_2	M_3	M_4	M_5
$ V_k $	$\frac{3}{2}(3^{k-1} + 1)$	$2(4^{k-1} + 1)$	$2(4^{k-1} + 1)$	$2(4^{k-1} + 1)$	$2(4^{k-1} + 1)$
$ E_k $	3^k	4^k	4^k	$5 \cdot 4^{k-1}$	$6 \cdot 4^{k-1}$
	$\frac{3}{2}(3^{k-1} - 1)p$	$\frac{4}{3}(4^{k-1} - 1)p$	$\frac{4}{3}(4^{k-1} - 1)p$	$\frac{5}{3}(4^{k-1} - 1)p$	$2(4^{k-1} - 1)p$
$\langle n_k \rangle$	$4 + 2p$	$4 + \frac{4}{3}p$	$4 + \frac{4}{3}p$	$5 + \frac{5}{3}p$	$6 + 2p$
	$-4 \frac{1+p}{3^{k-1}+1}$	$-\frac{4}{3} \frac{3+2p}{4^{k-1}+1}$	$-\frac{4}{3} \frac{3+2p}{4^{k-1}+1}$	$-\frac{5}{3} \frac{3+2p}{4^{k-1}+1}$	$-2 \frac{3+2p}{4^{k-1}+1}$
$\text{diam}(\Lambda_k), p = 0$	2^{k-1}	2^k	2^k	2^k	2^{k-1}
$\text{diam}(\Lambda_k), p = 1$	k	$k + 1$	$2(k - 1)$	k	k
$C, p = 1$	$(\log_6 3)^{-1}$	$(\log_6 4)^{-1}$	$2(\log_6 4)^{-1}$	$(\log_7 4)^{-1}$	$(\log_8 4)^{-1}$

In table 1, we collected the structural characteristics of our graphs based on the motifs M_1, \dots, M_5 given in figure 1. Its first three rows are obtained from the formulas (9) and (11).

3. Graph structure and the Ising model properties

As was mentioned above, there exists a profound connection between the properties of Gibbs random fields of models based on graphs and the structural properties of these graphs⁴. For the hierarchical lattices, the notion of the Gibbs random field of the Ising model was introduced in [14, 15]. In the physical terminology, each (pure) Gibbs random field corresponds to a state of thermal equilibrium of the model, see [25] for more details. Accordingly, the existence of multiple Gibbs random fields corresponds to the existence of multiple equilibrium states and hence of phase transitions. If there is no interaction between the spins, the Gibbs random field is unique. However, if the interaction is strong enough and if it is effectively propagated by the underlying graph (high enough “connectivity”), the Gibbs fields can be multiple. The condition of high connectivity is essential, which can be seen from the fact that the Ising model on the one-dimensional lattice \mathbb{Z} , considered as a graph with the natural adjacency relation, bears only one Gibbs field and hence no phase transitions can occur in this case – no matter how strong the interaction is. With regard to these arguments, we can divide all graphs into three groups according to the following property of the Ising model defined on these graphs:

⁴Of course, we are speaking now about countably infinite graphs.

- (a) There exists only one Gibbs random field for all interactions.
- (b) There exists only one Gibbs random field for weak interactions, and multiple Gibbs random fields for strong interactions.
- (c) There exist multiple Gibbs random fields for all nonzero interactions.

As was mentioned above, the lattice \mathbb{Z} belongs to the first group. The lattices \mathbb{Z}^d with $d \geq 2$, as well as all Cayley trees⁵ belong to the second group. An example of the graph belonging to the third group can be found in section 4 of [26]. Clearly, the above classification should be made more precise for random graphs, since in this case the existence of a Gibbs random field is a random event. We return to this issue below.

The Ising model defined on the graphs from the second group may have a *critical point*, which separates the regime of uniqueness (weak interactions) from the regime of non-uniqueness (strong interactions). At their critical point, the Gibbs random fields have “unusual” properties, which can be detected without explicit construction of these fields. One of such properties is the so-called *self-similarity*, which manifests itself in the appearance of unstable fixed points of some (renormalization) transformations. It turns out that the hierarchical graphs are quite suitable for such a study — that is why they appear in this context. Thus, one can have the following criterion: if the Ising model has a critical point, the graph belongs to the second group. If not, then the graph belongs either to the first or to the third group. The latter cases can be distinguished by an additional study.

In contrast to the hierarchical lattices introduced and studied in [16], our bare graphs belong to the first group. This can be seen from the analysis which we present below — a direct proof of such a statement will be done in our forthcoming publication. Thus, the interaction along the decorating bonds plays the main role in the possible appearance of phase transitions in the Ising model defined on our graphs. In view of this fact, we will assume that the solid bonds bear the interaction K , whereas the dotted bonds bear the interaction L . As the bonds of the latter kind appear at random, we can think of the corresponding model as of the one defined on the fully decorated graph with random interactions along the decorating bonds. Thus, in this model, the interaction along each bond $\langle i, j \rangle$ is a random variable, denoted by L_{ij}^ω , which takes values 0 and some nonzero $L \in \mathbb{R}$ with probabilities $1-p$ and p , respectively. For different bonds, these random variables are independent.

For a given $k \in \mathbb{N}$, the Ising model on the fully decorated graph Λ_k is defined in the usual way by assigning spin variables $\sigma_i = \pm 1$ to the nodes $i \in V_k$ and by setting the Hamiltonian

$$-\beta\mathcal{H}_k = h \sum_{i \in V_k} \sigma_i + K \sum_{\langle i, j \rangle \in E'_k} \sigma_i \sigma_j + \sum_{\langle i, j \rangle \in E''_k} L_{ij}^\omega \sigma_i \sigma_j, \quad h, K \in \mathbb{R}, \quad k \in \mathbb{N}, \quad (28)$$

where h is an external field. In accordance with the above arguments, the third summand corresponds to the interaction along decorating bonds. We have included the first term in view of the following arguments. As is known, the Ising model on the lattices \mathbb{Z}^d , $d \geq 2$, exhibits phase transitions only if $h = 0$. This is also the case if the graph is amenable and quasi-transitive. For nonamenable graphs, this model may have a phase transition for nonzero h , see [27].

By the construction of our graphs, each Λ_k has q external nodes, i_1, \dots, i_q , and the corresponding number of internal nodes, see table 1. Such external nodes can be considered as a boundary of Λ_k . Correspondingly, we call the spin variables *external* or *boundary* (respectively, *internal*) spins if they are assigned to external (respectively, internal) nodes. For a fixed configuration of the external spins $\sigma_{i_1}, \dots, \sigma_{i_q}$, we consider

$$Z_k^\omega(\sigma_{i_1}, \dots, \sigma_{i_q}) = \sum_{\text{internal spins of } \Lambda_k} \exp(-\beta\mathcal{H}_k). \quad (29)$$

This is the conditional partition function of the spin system in Λ_k , conditioned by the fixed configuration of spins on the boundary of Λ_k . Of course, it depends on the graph, i.e. on the motif

⁵Except for \mathbb{Z} , see page 247 in [25].

used in its construction. According to the main principle of the theory of Gibbs random fields applied in our context, see [25], the fact that such a field is unique is ensured by the conditional partition function asymptotic independence, as $k \rightarrow +\infty$, of the boundary spins. For our graphs, Z_k^ω are random and hence also depend on ω . In this situation, one can apply the following two approaches: the above mentioned independence holds (i) for almost all ω ; (ii) in average. The latter corresponds to the fact that the expected values of Z_k^ω , which we denote by $\langle Z_k^\omega \rangle$, are independent of the boundary spins. These two approaches are called *quenched* and *annealed* disorders, respectively, see [28], page 99. We will work in the annealed approach. It can be shown that the summands in (29) are positive and separated from zero. Since the number of these summands increases to $+\infty$ as $k \rightarrow +\infty$, we have $\langle Z_k^\omega \rangle \rightarrow +\infty$. Therefore, a weaker form of the uniqueness condition can be

$$\lim_{k \rightarrow +\infty} \langle Z_k^\omega(\sigma_{i_1}, \dots, \sigma_{i_q}) \rangle / \langle Z_k^\omega(\sigma'_{i_1}, \dots, \sigma'_{i_q}) \rangle = c, \quad (30)$$

which holds for any two configurations of the boundary spins and for some $c > 0$, which depends on these configurations. The convergence in (30) should be stable with respect to small changes of h and K , i.e. of the starting element Z_1 . In this case, the annealed limiting free energy per spin is independent of the boundary configuration. Correspondingly, a necessary condition for the latter quantity to be dependent on the boundary spins is that

$$\lim_{k \rightarrow +\infty} \langle Z_k^\omega(\sigma_{i_1}, \dots, \sigma_{i_q}) \rangle / \langle Z_k^\omega(\sigma'_{i_1}, \dots, \sigma'_{i_q}) \rangle = +\infty, \quad (31)$$

for some $\sigma_{i_1}, \dots, \sigma_{i_q}$ and $\sigma'_{i_1}, \dots, \sigma'_{i_q}$. We say that the Ising model defined on our graph has a critical point, if there exists $c' > 0$, distinct from c in (30), such that for $K = K_*$ and $h = h_*$,

$$\lim_{k \rightarrow +\infty} \langle Z_k^\omega(\sigma_{i_1}, \dots, \sigma_{i_q}) \rangle / \langle Z_k^\omega(\sigma'_{i_1}, \dots, \sigma'_{i_q}) \rangle = c'. \quad (32)$$

Here K_* and h_* are certain values of the interaction intensity and the external field. The convergence in (32) should be such that for arbitrarily small deviations from the point (K_*, h_*) one has either (30) or (31) instead of (32). Note that we do not exclude that $h_* \neq 0$ since our graphs are not quasi-transitive and hence the result of [27] may not be applicable in this case.

4. Critical points of the Ising model on M_1 based graph

For $k = 1$, we have only basic bonds and no inner nodes. Thus,

$$Z_1(a, b, c) = \exp[-\beta \mathcal{H}_1(a, b, c)] = \exp[K(ab + ac + bc) + h(a + b + c)], \quad (33)$$

where we use the same notations for the nodes a, b, c , see the left-hand graph in figure 2, as well as for the corresponding spin variable, that is, for short we write $a = \sigma_a$, $b = \sigma_b$, ect. For $k = 2$, we have, see the middle graph in figure 2,

$$-\beta \mathcal{H}_2(a, b, c, \alpha, \beta, \gamma) = -\beta \mathcal{H}_1(a, \gamma, \beta) - \beta \mathcal{H}_1(\gamma, b, \alpha) - \beta \mathcal{H}_1(\beta, \alpha, c) + L_{ab}^\omega ab + L_{bc}^\omega bc + L_{ac}^\omega ac.$$

Thus, to obtain $Z_2(a, b, c)$ we have to sum out the internal spins, see (29). That is,

$$Z_2^\omega(a, b, c) = \exp[L_{ab}^\omega ab + L_{ac}^\omega ac + L_{bc}^\omega bc] \sum_{\alpha, \beta, \gamma} Z_1(a, \gamma, \beta) Z_1(\gamma, b, \alpha) Z_1(\beta, \alpha, c),$$

where the sum is taken over $\alpha, \beta, \gamma = \pm 1$. According to the hierarchical structure of the underlying graph, for all $k \geq 2$ the graph Λ_k is obtained from the subgraphs Λ_{k-1} exactly in the same way. This yields the following recurrence for the partition functions (29)

$$Z_k^\omega(a, b, c) = R_k^\omega(a, b, c) \sum_{\alpha, \beta, \gamma} Z_{k-1}(a, \gamma, \beta) Z_{k-1}(\gamma, b, \alpha) Z_{k-1}(\beta, \alpha, c), \quad (34)$$

where

$$R_k^\omega(a, b, c) = \exp [L_{ab}^\omega ab + L_{ac}^\omega ac + L_{bc}^\omega bc], \quad (35)$$

and $Z_1(a, b, c)$ is given in (33). From the independence of L_{ij}^ω for different bonds, it follows that the random variables Z_{k-1}^ω on the right-hand side of (34) are independent, which allows us to compute the expectations and obtain

$$\langle Z_k^\omega(a, b, c) \rangle = \langle R_k^\omega(a, b, c) \rangle \sum_{\alpha, \beta, \gamma} \langle Z_{k-1}^\omega(a, \beta, \gamma) \rangle \langle Z_{k-1}^\omega(\alpha, b, \gamma) \rangle \langle Z_{k-1}^\omega(\alpha, \beta, c) \rangle. \quad (36)$$

We recall that the random variables L_{ij}^ω take the values $L \neq 0$ and 0 with probabilities p and $1 - p$, respectively. In view of the model symmetry, c.f. (33), the system (36) involves the following variables

$$\begin{aligned} A_k &= \langle Z_k^\omega(1, 1, 1) \rangle, & B_k &= \langle Z_k^\omega(1, 1, -1) \rangle, \\ C_k &= \langle Z_k^\omega(-1, -1, 1) \rangle, & D_k &= \langle Z_k^\omega(-1, -1, -1) \rangle, \end{aligned} \quad (37)$$

which have the initial values

$$A_1 = e^{3(K+h)}, \quad B_1 = e^{-K+h}, \quad C_1 = e^{-K-h}, \quad D_1 = e^{3(K-h)}. \quad (38)$$

By direct calculations,

$$\begin{aligned} \langle R_k^\omega(1, 1, 1) \rangle &= (p \exp(L) + 1 - p)^3, \\ \langle R_k^\omega(-1, 1, 1) \rangle &= (p \exp(L) + 1 - p)(p \exp(-L) + 1 - p)^2. \end{aligned} \quad (39)$$

Thereafter, the recursion (36) can be written in the form

$$\begin{aligned} x_{k+1} &= t \frac{P_x(x_k, y_k)}{Q(x_k, y_k, z_k)}, \\ y_{k+1} &= \frac{P_y(x_k, y_k, z_k)}{Q(x_k, y_k, z_k)}, \\ z_{k+1} &= t \frac{P_z(y_k, z_k)}{Q(x_k, y_k, z_k)}, \end{aligned} \quad (40)$$

with the initial conditions

$$x_1 = e^{4(K+h)}, \quad y_1 = e^{2h}, \quad z_1 = e^{4K-2h}, \quad (41)$$

where we have used the notations

$$x_k = A_k/C_k, \quad y_k = B_k/C_k, \quad z_k = D_k/C_k, \quad k \in \mathbb{N}, \quad (42)$$

and

$$t = \left(\frac{p e^L + 1 - p}{p e^{-L} + 1 - p} \right)^2. \quad (43)$$

The polynomials in (40) have the following form

$$\begin{aligned} P_x(x, y) &= x^3 + 3xy^2 + 3y^2 + 1, \\ P_y(x, y, z) &= x^2y + y^3 + 2xy + y^2z + 2y + z, \\ P_z(y, z) &= y^3 + 3y + 3z + z^3, \\ Q(x, y, z) &= xy^2 + x + 2y^2 + 2yz + z^2 + 1. \end{aligned} \quad (44)$$

For every $t > 0$, the mapping $(x_k, y_k, z_k) \mapsto (x_{k+1}, y_{k+1}, z_{k+1}) = T_t(x_k, y_k, z_k)$, defined in (40) and (44), maps the first octant of \mathbb{R}^3 into itself. The starting point (x_1, y_1, z_1) lies on the surface \mathcal{S} defined by the equation, c.f. (41),

$$x = y^3 z, \quad (45)$$

which, however, is not preserved by this mapping, which can be checked directly. Let \mathcal{T}_t be the invariant set of the map T_t , that is, the set of points (x, y, z) such that $T_t(x, y, z) = (x, y, z)$. Suppose that the intersection of \mathcal{T}_t with \mathcal{S} is non-void. If (x_*, y_*, z_*) belongs to this intersection and $(x_1, y_1, z_1) = (x_*, y_*, z_*)$, then $(x_k, y_k, z_k) = (x_*, y_*, z_*)$ for all $k \in \mathbb{N}$. Let us find such points. From the second equation in (40) for $x = y^3 z$ we get

$$yQ(y^3 z, y, z) = P_y(y^3 z, y, z),$$

which can be transformed into the following

$$y^3(1 + y^3 z)(1 - yz) = (y + z)(1 - yz). \quad (46)$$

A solution of the latter equation is $z = 1/y$, thus $x = y^2$, which being used in the first equation in (40) yields $(1 + y^2)^2 = t(1 + y^2)^2$ and hence $t = 1$. For $p > 0$, the latter implies $L = 0$, and also $K = 0$, see (41). This fixed point corresponds to a noninteracting system of spins in an external field and thereby has nothing to do with critical points. For $z \neq 1/y$, from (46) we get

$$z(1 - y^2)(y^4 + y^2 + 1) = -y(1 - y^2), \quad (47)$$

which for positive y and z has a unique solution $y = 1$, and hence $x = z$. We insert this into the first equation in (40) and obtain

$$x = t \frac{x^3 + 3x + 4}{x^2 + 4x + 3} = t \frac{x^2 - x + 4}{x + 3} := \phi_t(x). \quad (48)$$

For $t = 1$, the only solution is $x = 1$, which corresponds to $L = K = 0$. The choice $t = 1$ also corresponds to a bare graph ($p = 0$); hence, the bare graph based on M_1 has no critical points and no phase transitions. As we have already mentioned above, in a separate work we prove that the Gibbs states of the Ising model based on this graph are always unique. For $t < 1$, that is, for $L < 0$, (48) has only one positive solution

$$x_* = \frac{-3 - t + \sqrt{9 + 22t - 15t^2}}{2(1 - t)} < 1, \quad (49)$$

which corresponds to $K_* < 0$. It is stable and $(x_k, y_k, z_k) \rightarrow (x_*, 1, x_*)$ as $k \rightarrow +\infty$, for any $(x_1, y_1, z_1) \in \mathcal{S}$. Thus, for $L < 0$, the Ising model has no critical points and no phase transitions, even for the fully decorated graph. This can be caused by a frustration, which takes place in an antiferromagnetic Ising model on triangles. For $t > 1$, figure 5 presents the graphical solutions of (48). They have the following form

$$x_*^{(1)} = \frac{3 + t - \sqrt{9 + 22t - 15t^2}}{2(t - 1)}, \quad x_*^{(2)} = \frac{3 + t + \sqrt{9 + 22t - 15t^2}}{2(t - 1)}. \quad (50)$$

By direct calculations, we get $\phi'_t(x_*^{(1)}) < 1$ and $\phi'_t(x_*^{(2)}) > 1$, see figure 5. Hence, $x_*^{(1)}$ is stable, whereas $x_*^{(2)}$ is unstable. Note also that $x_*^{(1)} \rightarrow 1$ and $x_*^{(2)} \rightarrow \infty$ as $t \rightarrow 1$. The solutions (50) exist and are considered to be distinct, provided $t \in (1, 9/5)$. This yields the condition

$$p < \psi(L) := \frac{3 - \sqrt{5}}{\sqrt{5} \exp(L) - 3 \exp(-L) + 3 - \sqrt{5}}. \quad (51)$$

Since $\psi(L)$ is a decreasing function, the equation $\psi(L) = 1$ has only one solution

$$L_* = \frac{1}{4} \ln \frac{9}{5}. \quad (52)$$




Figure 5. Graphical solution of (48).

Then, for $L \leq L_*$, the critical point exists for all $p \in (0, 1]$. For such L and p , let $K_*(L, p)$ be the solution of the equation

$$\exp(4K) = x_*^{(2)}. \quad (53)$$

Then

$$\lim_{k \rightarrow +\infty} x_k = \begin{cases} x_*^{(1)} & \text{if } x_1 < \exp[4K_*(L, p)], \\ x_*^{(2)} & \text{if } x_1 = \exp[4K_*(L, p)], \\ +\infty & \text{if } x_1 > \exp[4K_*(L, p)]. \end{cases} \quad (54)$$

The conditions in (54) can be formulated directly for K , e.g. $x_k \rightarrow x_*^{(1)}$ if $K < K_*(L, p)$. Thus, $x_*^{(1)}$ is the so-called high-temperature fixed point. For $L > L_*$, we have $\psi(L) < 1$. Hence, the condition (51) is not satisfied for $p \in (\psi(L), 1]$, which also includes the fully decorated graph with $p = 1$. For such L and p , the whole graph of ϕ_t lies above the line $\phi_t(x) = x$, which means that $x_k \rightarrow +\infty$ for all initial $x_1 \geq 0$. Thus, for $L > L_*$ and $p \in (\psi(L_*), 1]$, the model has no critical points and the Gibbs random fields are multiple for all $K \in \mathbb{R}$. For $p = \psi(L_*)$, we have $t = 9/5$ and $x_*^{(1)} = x_*^{(2)} = 3$, which corresponds to $K_* = (\ln 3)/4$. In this case, $x_k \rightarrow 3$ if $K \leq K_*$, and $x_k \rightarrow +\infty$ if $K > K_*$.

5. Concluding remarks

5.1. Construction and structural properties

The graphs introduced in this paper are constructed from motifs M_1, \dots, M_5 in an algorithmic way, like the hierarchical lattices introduced in [13], and employed in [16] where they were supplied with long-range (decorating) bonds. However, our main construction principle — to replace nodes with the graphs of the previous level — differs from the one used in those papers, where the graphs of the previous level replaced the bonds. Hierarchical graphs of this kind model the real world networks with the so-called modular structure, which are organized in tightly knit communities with relatively sparse connections between them. For more details herein, we refer the reader to [29, 30] and to the publications cited therein. In contrast to the graphs introduced in [16], see also [29], our decorated graphs are not scale-free. For them, the node degree distribution is of intermediate type such that all the moments exist. The corresponding characteristic function can be extended to a meromorphic function, analytic in the complex neighborhood of zero, see (18), whereas for scale-free graphs such functions are nonanalytic at zero. All our graphs are amenable, which correlates with the property just discussed, see (14), (9), and (19). An interesting property is that, like for the graphs studied in [16], adding decorations forms the graphs possessing the small-world property.

5.2. Critical points

The appearance of critical points, and hence of phase transitions, for the Ising model confirms good communicating properties of the underlying graphs. For our graphs, in contrast to the hierarchical lattices for which phase transitions take place even without decorations, critical points can exist only if the decorations are applied, with any $p > 0$. The most interesting related fact is that for the graph based on M_1 , the antiferromagnetic Ising model has no phase transitions for any L and p . This is due to the generic frustration of such models — the elementary factor $R_k(a, b, c)$ which appears in (35) with negative L cannot be maximized. We expect that the same is true for the graphs based on M_5 . To check if this conjecture of ours is true we plan to study a Potts model on the same graph and $q \geq 3$, for which such a frustration is no longer actual. We also plan to look for critical points with nonzero h , as well as for critical points of the Ising model with the underlying graphs based on the remaining motifs. However, the corresponding problems are much tougher, so that the appropriate numerical means should be applied.

6. Acknowledgement

The authors benefited from fruitful discussions on the matter of this work held with our colleagues Yuri Kondratiev and Vasyl Ustimenko for which they are cordially grateful. The authors are also grateful to the unnamed referee whose remarks and suggestions were helpful in improving the presentation of the article.

References

1. Erdős P., Rényi A., *Publicationes Mathematicae*, 1959, **6**, 290.
2. Erdős P., Rényi A., *Publications of the Mathematical Institute of the Hungarian Academy of Sciences*, 1960, **5**, 17.
3. Białas, P., Burda, Z., Waclaw, B. Causal and homogeneous networks. In: *Science of complex networks*, 14, AIP Conf. Proc., **776**, Amer. Inst. Phys., Melville, NY, 2005.
4. Watts D.J., Strogatz H., *Nature*, 1998, **393**, 440; doi:10.1038/30918.
5. Barabási A.-L., Albert R., *Science*, 1999, **286**, 509; doi:10.1126/science.286.5439.509.
6. Milo R. et al., *Science*, 2002, **298**, 824; doi:10.1126/science.298.5594.824.
7. Itzkovitz S. et al., *Phys. Rev. E*, 2003, **68**, 026127; doi:10.1103/PhysRevE.68.026127.
8. Alon U., *Science*, 2003, **301**, 1866; doi:10.1126/science.1089072; *Nature Reviews Genetics*, 2007, **8**, 450; doi:10.1038/nrg2102.
9. Kashtan N., Itzkovitz S., Milo R., Alon U., *Phys. Rev. E*, 2004, **70**, 031909; doi:10.1103/PhysRevE.70.031909.
10. Itzkovitz S., Alon U., *Phys. Rev. E*, 2005, **71**, 026117; doi:10.1103/PhysRevE.71.026117.
11. Matias C. et al, *REVSTAT*, 2006, **4**, 31.
12. Häggström O., *Adv. Appl. Probab.*, 2000, **32**, 39.
13. Griffiths R.B., Kaufman M., *Phys. Rev. B*, 1982, **26**, 5022; doi:10.1103/PhysRevB.26.5022.
14. Bleher P.M., Žalys E., *Litovsk. Mat. Sb.*, 1988, **28**, 252 [English translation in *Lithuanian Math. J.*, 1989, **28**, 127; doi:10.1007/BF01027189].
15. Bleher P.M., Žalys E., *Commun. Math. Phys.*, 1989, **120**, 409; doi:10.1007/BF01225505.
16. Hinczewski M., A. Nihat Berker, *Phys. Rev. E*, 2006, **73**, 066126; doi:10.1103/PhysRevE.73.066126.
17. Wróbel M., *Condens. Matter Phys.*, 2008, **11**, 341.
18. Kotorowicz M., *Albanian Journal of Mathematics*, 2008, **3**, 235.
19. Barlow M.T. Heat kernels and sets with fractal structure. In: *Heat kernels and analysis on manifolds, graphs, and metric spaces* (Paris, 2002). *Contemp. Math.*, 338, Amer. Math. Soc., Providence, RI, 2003.
20. Newman M.E.J., *SIAM Review*, 2003, **45**, 167; doi:10.1137/S003614450342480.
21. Ruelle D., *Statistical mechanics: Rigorous results*. W.A. Benjamin, Inc., New York–Amsterdam, 1969.
22. Erdős P., Kleitman D.J., Rothschild B.L., Asymptotic enumeration of K_n -free graphs. In: *Colloquio Internazionale sulle Teorie Combinatorie* (Rome, 1973), Tomo II, pp. 19. *Atti dei Convegni Lincei*, No. 17, Accad. Naz. Lincei, Rome, 1976.
23. Futorny V., Ustimenko V., *Acta Appl. Math.*, 2007, **98**, 47; doi:10.1007/s10440-007-9144-8.

24. Barrat A., Weigh M., Eur. Phys. J. B, 2000, **13**, 547; doi:10.1007/s100510050067.
25. Georgii H.-O., Gibbs measures and phase transitions. de Gruyter Studies in Mathematics, **9**. Walter de Gruyter & Co., Berlin, 1988.
26. Kępa D., Kozitsky Y., Condens. Matter Phys., 2008, **11**, 313.
27. Jonasson J, Steif J.E., J. Theoret. Probab., 1999, **12**, 549; doi:10.1023/A:1021690414168.
28. Bovier, A., Statistical mechanics of disordered systems. A mathematical perspective. Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, Cambridge, 2006; doi:10.1017/CBO9780511616808.004.
29. Hinczewski M., Phys. Rev. E, 2007, **75**, 06611104; doi:10.1103/PhysRevE.75.061104.
30. Kashtan N., Mayo A.E., Kalisky T., Alon U., PLoS Comput. Biol., 2009, **5**, e1000355; doi:10.1371/journal.pcbi.1000355.

Ієрархічні випадкові графи з мотивів: структурні властивості та критичні точки моделі Ізінга

М. Которович, Юрій Козицький

Інститут математики, Університет Марії Кюрі-Склодовської, Люблін

Вводиться і вивчається клас випадкових графів, збудованих в алгоритмічний спосіб з п'яти мотивів, знайдених у [Milo R., Shen-Orr S., Itzkovitz S., Kashtan N., Chklovskii D., Alon U., Science, 2002, **298**, 824]. Конструкційна схема нагадує схему, застосовану у [Hinczewski M., A. Nihat Berker, Phys. Rev. E, 2006, **73**, 066126], згідно з якою короткосяжні ребра є невідповідні, тоді як довгосяжні ребра виникають незалежно і з однаковою ймовірністю. Описано ряд структурних властивостей графів, серед яких є розподіл ступенів, кластерність, аменабільність, властивість тісного світу. Для одного з мотивів вивчається критична точка моделі Ізінга, визначеної на відповідному графі.

Ключові слова: аменабільність, розподіл ступенів, кластерність, граф тісного світу, модель Ізінга, критична точка