

Mathematical Physics

KP and Toda tau functions in Bethe ansatz

Kanehisa Takasaki

(Submitted on 16 Mar 2010)

Recent work of Foda and his group on a connection between classical integrable hierarchies (the KP and 2D Toda hierarchies) and some quantum integrable systems (the 6-vertex model with DWBC, the finite XXZ chain of spin 1/2, the phase model on a finite chain, etc.) is reviewed. Some additional information on this issue is also presented.

Comments: latex2e, using ws-procs9x6 package, 19 pages, contribution to the festschrift volume for the 60th anniversary of Tetsuji Miwa

Subjects: **Mathematical Physics (math-ph)**; High Energy Physics - Theory (hep-th); Quantum Algebra (math.QA); Exactly Solvable and Integrable Systems (nlin.SI)

Cite as: [arXiv:1003.3071v1](#) [math-ph]

Submission history

From: Kanehisa Takasaki [[view email](#)]

[v1] Tue, 16 Mar 2010 03:51:10 GMT (23kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math-ph

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1003](#)

Change to browse by:

[hep-th](#)

[math](#)

[math.QA](#)

[nlin](#)

[nlin.SI](#)

References & Citations

- [CiteBase](#)

Bookmark (what is this?)

[CiteULike logo](#)

[Connotea logo](#)

[BibSonomy logo](#)

[Mendeley logo](#)

[Facebook logo](#)

[del.icio.us logo](#)

[Digg logo](#)

[Reddit logo](#)