

Quantum Physics

A glance at singlet states and four-partite correlations

Maria Schimpf, Karl Svozil

(Submitted on 7 Jan 2009 (v1), last revised 29 Jan 2009 (this version, v3))

Group theoretic methods to construct all N-particle singlet states by iterative recursion are presented. These techniques are applied to the quantum correlations of four spin one-half particles in their singlet states. Multipartite quantized systems can be partitioned, and their observables grouped and redefined into condensed correlations.

Comments: 24 pages, 7 figures, to appear in a special issue of Mathematica Slovaca dedicated to Sylvia Pulmannova

Subjects: **Quantum Physics (quant-ph)**

Cite as: **arXiv:0901.0882v3 [quant-ph]**

Submission history

From: Svozil Karl [[view email](#)]

[v1] Wed, 7 Jan 2009 19:29:02 GMT (452kb)

[v2] Thu, 8 Jan 2009 13:58:54 GMT (454kb)

[v3] Thu, 29 Jan 2009 12:59:09 GMT (455kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

quant-ph

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [0901](#)

References & Citations

- [SLAC-SPIRES HEP](#)
([refers to](#) | [cited by](#))
- [CiteBase](#)

Bookmark (what is this?)

[CiteULike logo](#)

[Connotea logo](#)

[BibSonomy logo](#)

[Mendeley logo](#)

[Facebook logo](#)

[del.icio.us logo](#)

[Digg logo](#)

[Reddit logo](#)