

Quantum Physics

Semidefinite programs for completely bounded norms

[John Watrous](#)

(Submitted on 29 Jan 2009 (v1), last revised 15 Apr 2009 (this version, v2))

The completely bounded trace and spectral norms in finite dimensions are shown to be expressible by semidefinite programs. This provides an efficient method by which these norms may be both calculated and verified, and gives alternate proofs of some known facts about them.

Comments: 17 pages, references added in version 2

Subjects: **Quantum Physics (quant-ph)**

Cite as: **arXiv:0901.4709v2 [quant-ph]**

Submission history

From: John Watrous [[view email](#)]

[v1] Thu, 29 Jan 2009 15:53:13 GMT (16kb)

[v2] Wed, 15 Apr 2009 17:39:45 GMT (17kb)

[Which authors of this paper are endorsers?](#)

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

quant-ph

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [0901](#)

References & Citations

- [SLAC-SPIRES HEP](#)
([refers to](#) | [cited by](#))
- [CiteBase](#)

Bookmark (what is this?)

[CiteULike logo](#)

[Connotea logo](#)

[BibSonomy logo](#)

[Mendeley logo](#)

[Facebook logo](#)

[del.icio.us logo](#)

[Digg logo](#)

[Reddit logo](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).