Quantum Physics

Quantum phase diagram of fermion mixtures with population imbalance in one-dimensional optical lattices

B. Wang, Han-Dong Chen, S. Das Sarma

(Submitted on 30 Jan 2009)

With a recently developed time evolving block decimation (TEBD) algorithm, we numerically study the ground state quantum phase diagram of fermi mixtures with attractive inter-species interactions loaded in one-dimensional optical lattices. For our study, we adopt a general asymmetric Hubbard model (AHM) with species-dependent tunneling rates to incorporate the possibility of mass imbalance in the mixtures. We find clear signatures for the existence of a Fulde-Ferrell-Larkin-Ovchinnikov (FFLO) phase in this model in the presence of population imbalance. Our simulation also reveals that in the presence of mass imbalance, the parameter space for FFLO states shrinks or even completely vanishes depending on the strength of the attractive interaction and the degree of mass imbalance.

Subjects:Quantum Physics (quant-ph); Strongly Correlated Electrons
(cond-mat.str-el)Journal reference:Phys. Rev. A 79, 051604(R) (2009)DOI:10.1103/PhysRevA.79.051604Cite as:arXiv:0901.4896v1 [quant-ph]

Submission history

From: Bin Wang [view email] [v1] Fri, 30 Jan 2009 14:46:27 GMT (744kb)

Which authors of this paper are endorsers?

Link back to: arXiv, form interface, contact.

(Help | Advanced search)

All papers - Go!

Download:

- PDF
- PostScript
- Other formats

Current browse context: quant-ph < prev | next > new | recent | 0901

Change to browse by:

cond-mat cond-mat.str-el

References & Citations

- SLAC-SPIRES HEP (refers to | cited by)
- CiteBase

Bookmark(what is this?)

