

# Stationary states and fractional dynamics in systems with long range interactions

Tineke L. Van Den Berg (IM2NP), Duccio Fanelli, Xavier Leoncini (CPT)

(Submitted on 16 Dec 2009)

Dynamics of many-body Hamiltonian systems with long range interactions is studied, in the context of the so called  $\alpha$ -HMF model. Building on the analogy with the related mean field model, we construct stationary states of the  $\alpha$ -HMF model for which the spatial organization satisfies a fractional equation. At variance, the microscopic dynamics turns out to be regular and explicitly known. As a consequence, dynamical regularity is achieved at the price of strong spatial complexity, namely a microscopic inhomogeneity which locally displays scale invariance.

Subjects: **Statistical Mechanics (cond-mat.stat-mech)**; Chaotic Dynamics (nlin.CD)

Cite as: [arXiv:0912.3060v1](https://arxiv.org/abs/0912.3060v1) [cond-mat.stat-mech]

## Submission history

From: Xavier Leoncini [[view email](#)] [via CCSD proxy]  
[v1] Wed, 16 Dec 2009 07:01:14 GMT (286kb)

*[Which authors of this paper are endorsers?](#)*

## Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

cond-mat.stat-mech

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [0912](#)

Change to browse by:

[cond-mat](#)

[nlin](#)

[nlin.CD](#)

## References & Citations

- [CiteBase](#)

## Bookmark([what is this?](#))

[CiteULike logo](#)

[Connotea logo](#)

[BibSonomy logo](#)

[Mendeley logo](#)

[Facebook logo](#)

[del.icio.us logo](#)

[Digg logo](#)

[Reddit logo](#)