



Mathematical Physics

Green's function for the wavized Maxwell fish-eye problem

Radosław Szmytkowski

(Submitted on 7 Jul 2011)

Unique transformation properties under the hyperspherical inversion of a partial differential equation describing a stationary scalar wave in an N -dimensional ($N \geq 2$) Maxwell fish-eye medium are exploited to construct a closed form of the Green's function for that equation. For those wave numbers for which the Green's function fails to exist, the generalized Green's function is derived. Prospective physical applications are mentioned.

Comments: LaTeX, 14 pages
 Subjects: **Mathematical Physics (math-ph)**; Analysis of PDEs (math.AP); Optics (physics.optics)
 MSC classes: 35J08, 78A10
 Journal reference: J. Phys. A 44 (2011) 065203
 DOI: [10.1088/1751-8113/44/6/065203](https://doi.org/10.1088/1751-8113/44/6/065203)
 Cite as: [arXiv:1107.1466](https://arxiv.org/abs/1107.1466) [math-ph]
 (or [arXiv:1107.1466v1](https://arxiv.org/abs/1107.1466v1) [math-ph] for this version)

Submission history

From: Radoslaw Szmytkowski [[view email](#)]
 [v1] Thu, 7 Jul 2011 17:51:37 GMT (12kb)

Which authors of this paper are endorsers?

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

math-ph

< [prev](#) | [next](#) >

[new](#) | [recent](#) | [1107](#)

Change to browse by:

[math](#)

[math.AP](#)

[physics](#)

[physics.optics](#)

References & Citations

- [NASA ADS](#)

Bookmark (what is this?)

