

Vesicle electrohydrodynamics

Jonathan T. Schwalbe, Petia M. Vlahovska, Michael J. Miksis

(Submitted on 12 Nov 2010)

A small amplitude perturbation analysis is developed to describe the effect of a uniform electric field on the dynamics of a lipid bilayer vesicle in a simple shear flow. All media are treated as leaky dielectrics and fluid motion is described by the Stokes equations. The instantaneous vesicle shape is obtained by balancing electric, hydrodynamic, bending, and tension stresses exerted on the membrane. We find that in the absence of ambient shear flow, it is possible that an applied step--wise uniform DC electric field could cause the vesicle shape to evolve from oblate to prolate over time if the encapsulated fluid is less conducting than the suspending fluid. For a vesicle in ambient shear flow, the electric field damps the tumbling motion leading to a stable tank-treading state.

Comments: 15 pages, 7 figures

Subjects: **Fluid Dynamics (physics.flu-dyn)**

Cite as: **arXiv:1011.2995v1** [physics.flu-dyn]

Submission history

From: Jonathan Schwalbe [view email]

[v1] Fri, 12 Nov 2010 17:54:06 GMT (1483kb,D)

Which authors of this paper are endorsers?

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [Other formats](#)

Current browse context:

physics.flu-dyn

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1011](#)

Change to browse by:

[physics](#)

References & Citations

- [NASA ADS](#)

Bookmark([what is this?](#))

