

Nuclear Theory

Anisotropic collective flow of a Lorentz gas

Nicolas Borghini, Clement Gombeaud

(Submitted on 29 Jun 2011 (v1), last revised 30 Jun 2011 (this version, v2))

Analytical results for the anisotropic collective flow of a Lorentz gas of massless particles scattering on fixed centres are presented.

Comments: 8 pages, 1 figure. This is an extended version of what is being submitted (4 pages) to the proceedings of Quark Matter 2011, Annecy, May 22-28. v2: typos corrected

Subjects: **Nuclear Theory (nucl-th)**

Report number: BI-TP 2011/17

Cite as: [arXiv:1106.6009](https://arxiv.org/abs/1106.6009) [nucl-th]
(or [arXiv:1106.6009v2](https://arxiv.org/abs/1106.6009v2) [nucl-th] for this version)

Submission history

From: Nicolas Borghini [[view email](#)]

[v1] Wed, 29 Jun 2011 17:25:31 GMT (12kb)

[v2] Thu, 30 Jun 2011 14:32:36 GMT (12kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse contents:

nucl-th

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

References & Citations:

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

