

## High Energy Physics - Phenomenology

# Mixing angle between $\chi_{3P_1}$ and $\chi_{1P_1}$ in HQET

Takayuki Matsuki, Toshiyuki Morii, Kohichi Seo

*(Submitted on 24 Jan 2010)*

Some claim that there are two independent mixing angles ( $\theta = 35.3^\circ$ ,  $54.7^\circ$ ) between  $\chi_{3P_1}$  and  $\chi_{1P_1}$  states of heavy-light mesons in heavy quark symmetric limit, and others claim there is only one ( $\theta = 35.3^\circ$ ). We clarify the difference between these two and suggest which should be adopted. General arguments on the mixing angle between  $\chi_{3L_L}$  and  $\chi_{1L_L}$  of heavy-light mesons are given in HQET and a general relation is derived in heavy quark mass limit as well as that including the first order correction in  $1/m_Q$ .

Comments: 6 pages, 1 table

Subjects: **High Energy Physics - Phenomenology (hep-ph)**; High Energy Physics - Experiment (hep-ex); Nuclear Theory (nucl-th)Cite as: [arXiv:1001.4248v1](#) [hep-ph]

## Submission history

From: Takayuki Matsuki [[view email](#)]**[v1]** Sun, 24 Jan 2010 15:31:50 GMT (9kb)*[Which authors of this paper are endorsers?](#)*

## Download:

- [PostScript](#)
- [PDF](#)
- [Other formats](#)

Current browse context:

hep-ph

[< prev](#) | [next >](#)[new](#) | [recent](#) | [1001](#)

Change to browse by:

[hep-ex](#)[nucl-th](#)

## References & Citations

- [SLAC-SPIRES HEP](#)  
([refers to](#) | [cited by](#))
- [CiteBase](#)

Bookmark([what is this?](#))Link back to: [arXiv](#), [form interface](#), [contact](#).