

Plenary talk

Strategies for baryon resonance analysis

M. Döring¹, C. Hanhart^{1,2}, 黄飞³, S. Krewald^{1,2}, U. G. Meißner^{1,2,4}

1 Institut für Kernphysik and Jülich Center for Hadron Physics, Forschungszentrum Jülich, D-52425 Jülich, Germany

2 Institute for Advanced Simulation, Forschungszentrum Jülich, D-52425 Jülich, Germany

3 Department of Physics and Astronomy, University of Georgia, Athens, Georgia 30602, USA

4 Helmholtz-Institut für Strahlen- und Kernphysik (Theorie) and Bethe Center for Theoretical Physics, Universität Bonn, Nuß allee 14-16, D-53115 Bonn, Germany

收稿日期 2009-8-7 修回日期 网络版发布日期 2009-11-11 接受日期 2009-11-11

摘要

The analytic properties of scattering amplitudes provide a meeting point for experimental and theoretical investigations of baryon resonances. Pole positions and residues allow for a parameterization of resonances in a well-defined way which relates different reactions. The recent progress made within the Jülich model is summarized.

关键词 [pion-nucleon scattering, meson exchange model, analytic structure of the scattering amplitude](#)

分类号

DOI:

通讯作者:

M. Döring

作者个人主页:

M. Döring¹; C. Hanhart^{1,2}; 黄飞³; S. Krewald^{1,2}; U. G. Meißner^{1,2,4}

扩展功能

本文信息

► [Supporting info](#)

► [PDF](#)(735KB)

► [\[HTML全文\]\(0KB\)](#)

► [参考文献\[PDF\]](#)

► [参考文献](#)

服务与反馈

► [把本文推荐给朋友](#)

► [加入我的书架](#)

► [加入引用管理器](#)

► [引用本文](#)

► [Email Alert](#)

相关信息

► [本刊中包含“pion-nucleon scattering, meson exchange model, analytic structure of the scattering amplitude”的相关文章](#)

► 本文作者相关文章

· [M Douml](#)

· [ring](#)

· [C Hanhart](#)

·

· [黄飞](#)

· [S Krewald](#)

·

· [U G Meisslig](#)

· [ner](#)

·