

粒子天体物理与宇宙学

Plasma neutrino energy loss due to the axial-vector current at the late stages of stellar evolution

刘晶晶

(Department of Physics, Qiongzhou University, Wuzhishan 572200, China)

收稿日期 2008-10-20 修回日期 2008-12-20 网络版发布日期 2009-9-11 接受日期 2009-9-11

摘要

Based on the Weinberg-Salam theory, the plasma neutrino energy loss rates of vector and axial-vector contributions are studied. A ratable factor of the rates from the axial-vector current relative to those of the total neutrino energy loss rates is accurately calculated. The results show that the ratable factor will reach a maximum of 0.95 or even more at relatively higher temperature and lower density (such as  $\rho\mu_e < 10^7 \text{ g/cm}^3$ ). Thus the rates of the axial-vector contribution cannot be neglected. On the other hand, the rates of the axial-vector contribution are on the order of  $\sim 0.01\%$  of the total vector contribution, which is in good agreement with Itoh's at relatively high density (such as  $\rho\mu_e > 10^7 \text{ g/cm}^3$ ) and a temperature of  $T \leq 10^{11} \text{ K}$ .

关键词 [Weinberg-Salam theory, plasma neutrino energy loss, stellar evolution](#)

分类号

DOI:

通讯作者:

刘晶晶 [liujingjing68@126.com](mailto:liujingjing68@126.com)

作者个人主页:

刘晶晶

扩展功能

本文信息

▶ [Supporting info](#)

▶ [PDF](#) (1185KB)

▶ [\[HTML全文\]](#) (0KB)

▶ [参考文献\[PDF\]](#)

▶ [参考文献](#)

服务与反馈

▶ [把本文推荐给朋友](#)

▶ [加入我的书架](#)

▶ [加入引用管理器](#)

▶ [引用本文](#)

▶ [Email Alert](#)

相关信息

▶ [本刊中包含“Weinberg-Salam theory, plasma neutrino energy loss, stellar evolution”的 \[相关文章\]\(#\)](#)

▶ [本文作者相关文章](#)

· [刘晶晶](#)