



Nuclear Theory

Equation of state at finite baryon density based on lattice QCD

Pasi Huovinen, Peter Petreczky

(Submitted on 30 Jun 2011)

We employ the lattice QCD data on Taylor expansion coefficients to extend our previous parametrization of the equation of state to finite baryon density. When we take into account lattice spacing and quark mass dependence of the hadron masses, the Taylor coefficients at low temperature are equal to those of hadron resonance gas. Thus the equation of state is smoothly connected to the hadron resonance gas equation of state at low temperatures. We also show how the elliptic flow is affected by this equation of state at the maximum SPS energy.

Comments: 4 pages, 4 figures, Paraller talk at Quark Matter 2011, 22-28 May 2011, Anecy, France

Subjects: **Nuclear Theory (nucl-th)**; High Energy Physics - Lattice (hep-lat); High Energy Physics - Phenomenology (hep-ph); Nuclear Experiment (nucl-ex)

Cite as: [arXiv:1106.6227v1](https://arxiv.org/abs/1106.6227v1) [nucl-th]

Submission history

From: Pasi Huovinen [[view email](#)]

[v1] Thu, 30 Jun 2011 13:55:06 GMT (19kb)

[Which authors of this paper are endorsers?](#)

Link back to: [arXiv](#), [form interface](#), [contact](#).

Download:

- [PDF](#)
- [PostScript](#)
- [Other formats](#)

Current browse context:

nucl-th

[< prev](#) | [next >](#)

[new](#) | [recent](#) | [1106](#)

Change to browse by:

[hep-lat](#)

[hep-ph](#)

[nucl-ex](#)

References & Citations

- [INSPIRE HEP](#)
([refers to](#) | [cited by](#))
- [NASA ADS](#)

Bookmark([what is this?](#))

