

07 FORM FACTORS AND OZI RULE

Form factor measurements with PANDA at FAIR

E. Tomasi-Gustafsson (on behalf of the PANDA collaboration)

CEA, IRFU, SPhN, Saclay, 91191 Gif-sur-Yvette Cedex, France, and CNRS/IN2P3, Institut de Physique Nucléaire, UMR 8608, 91406 Orsay, France

收稿日期 2010-1-26 修回日期 网络版发布日期 2010-5-5 接受日期 2010-5-5

摘要

The high intensity and high energy antiproton beams which will be produced at FAIR open the possibility to determine time-like electromagnetic form factors in a wide kinematical range, through the annihilation reaction: $p^+p^- \rightarrow e^+e^-$. The status of the proposed experiment as well as the expected results are presented on the basis of realistic simulations. The impact of these measurements on the understanding of the nucleon structure, of the asymptotic properties of form factors and of the reaction mechanism are discussed using model independent statements based on symmetry properties of the strong and electromagnetic interactions in connection with space-like data.

关键词 [electromagnetic form factors, antiprotons, nucleon structure, QCD](#)

分类号

DOI:

通讯作者:

E. Tomasi-Gustafsson

作者个人主页:

E. Tomasi-Gustafsson (on behalf of the PANDA collaboration)

扩展功能

本文信息

▶ [Supporting info](#)

▶ [PDF \(1416KB\)](#)

▶ [\[HTML全文\]\(0KB\)](#)

▶ [参考文献\[PDF\]](#)

▶ [参考文献](#)

服务与反馈

▶ [把本文推荐给朋友](#)

▶ [加入我的书架](#)

▶ [加入引用管理器](#)

▶ [引用本文](#)

▶ [Email Alert](#)

相关信息

▶ [本刊中 包含“electromagnetic form factors, antiprotons, nucleon structure, QCD”的 相关文章](#)

▶ 本文作者相关文章

· [E Tomasi-Gustafsson on behalf of the PANDA collaboration](#)