

Turkish Journal of Chemistry

Turkish Journal

of

Chemistry

New Diterpenoids from the Aerial Parts of *Salvia pilifera*

Ufuk KOLAK

Department of General Chemistry, Faculty of Pharmacy, İstanbul

University, 34116, İstanbul-TURKEY

e-mail: ufukkolak@yahoo.com

 [Keywords](#)
 [Authors](#)

Abstract: Three new diterpenoids, piliferol (2α , 20-dihydroxy-12-oxo-abieta-9(11),13-dien-8,20-ether) (1), salvipiliferol (2α , 10β -dihydroxy-12-oxo-norabieta-9(11),13-dien) (2), and piliferalactone (2α , 8-dihydroxy-12-oxo-abieta-9(11),13-dien-20-oic acid-8,20-lactone) (3), together with 7 known compounds, ursolic acid, oleanolic acid, betulinic acid, α -amyrin, lupeol, β -sitosterol, and pectolarigenin, have been isolated from the aerial parts of an endemic *Salvia* species, *S. pilifera*. Their structures were elucidated by spectral (UV, IR, 1D-, and 2D-NMR, and HRMS) methods.

Key Words: *Salvia pilifera*, Lamiaceae, abietane-type diterpenoids, triterpenoids, steroid, flavone



chem@tubitak.gov.tr

[Scientific Journals Home Page](#)

Turk. J. Chem., **31**, (2007), 363-369.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Chem., vol.31, iss.3.](#)