

Turkish Journal of Chemistry

Turkish Journal

of

Chemistry

Terpenic and Phenolic Compounds from *Sideritis stricta*

F. Pinar ŞAHİN¹, Nurten EZER¹, İhsan ÇALIŞ²

¹Department of Pharmaceutical Botany, Faculty of Pharmacy, Hacettepe University,
TR-06100, Ankara-TURKEY

e-mail: psahin@hacettepe.edu.tr

²Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University,
TR-06100, Ankara-TURKEY

 [Keywords](#)
 [Authors](#)



chem@tubitak.gov.tr

[Scientific Journals Home](#)
[Page](#)

Abstract: From the overground parts of an endemic *Sideritis* species, *S. stricta*, 4 diterpenes, sideridiol (1), isosidol (2), isolinearol (3), and linearol (4); 2 flavonoid glycosides, isoscutellarein 7-O-[6"-O-acetyl-β-D-allopyranosyl-(1 to 2)]-β-D-glucopyranoside (5), and isoscutellarein 7-O-[6"-O-acetyl-β-D-allopyranosyl-(1 to 2)]-6"-O-acetyl-β-D-glucopyranoside (6); a methoxyflavone, xanthomicrol (7); and a phenylethanoid glycoside, verbascoside (=acteoside) (8) were isolated. The structures of the isolated compounds were established based on spectroscopic evidence (UV, IR, 1D- and 2D-NMR, MS).

Key Words: *Sideritis stricta*, diterpenoids, flavonoids, flavonoid glycosides, methoxyflavones, phenylethanoid glycosides

Turk. J. Chem., **30**, (2006), 495-504.

Full text: [pdf](#)

Other articles published in the same issue: [Turk. J. Chem.,vol.30,iss.4.](#)